PRODUCTS OF H -SEPARABLE SPACES IN THE LAVER MODEL

DUŠAN REPOVŠ AND LYUBOMYR ZDOMSKYY

ABSTRACT. We prove that in the Laver model for the consistency of the Borel's conjecture, the product of any two H -separable spaces is M -separable.

1. INTRODUCTION

This paper is devoted to products of H -separable spaces. A topological space X is said [3] to be H -separable, if for every sequence $\langle D_n : n \in \omega \rangle$ of dense subsets of X , one can pick finite subsets $F_n \subset D_n$ so that every nonempty open set $O \subset X$ meets all but finitely many F_n 's. If we only demand that $\bigcup_{n \in \omega} F_n$ is dense we get the definition of M -separable spaces introduced in [14]. It is obvious that second-countable spaces (even spaces with a countable π -base) are H -separable, and each H -separable space is M -separable. The main result of our paper is the following

Theorem 1.1. *In the Laver model for the consistency of the Borel's conjecture, the product of any two countable H -separable spaces is M -separable.*

Consequently, the product of any two H -separable spaces is M -separable provided that it is hereditarily separable.

It worth mentioning here that by [12, Theorem 1.2] the equality $\mathfrak{b} = \mathfrak{c}$ which holds in the Laver model implies that the M -separability is not preserved by finite products of countable spaces in the strong sense.

Let us recall that a topological space X is said to have the *Menger property* (or, alternatively, is a *Menger space*) if for every sequence $\langle \mathcal{U}_n : n \in \omega \rangle$ of open covers of X there exists a sequence $\langle \mathcal{V}_n : n \in \omega \rangle$ such that each $\mathcal{V}_n$ is a finite subfamily of $\mathcal{U}_n$ and the collection $\{\cup \mathcal{V}_n : n \in \omega\}$ is a cover of X . This property was introduced by Hurewicz, and the current name (the Menger property) is used because Hurewicz proved in [7] that for metrizable spaces his property is equivalent to a certain property of a base considered by Menger in [10]. If in the definition above we additionally require that $\{n \in \omega : x \notin \cup \mathcal{V}_n\}$ is finite for each $x \in X$, then we obtain the definition of the *Hurewicz property* introduced in [8]. The original idea behind the

2010 *Mathematics Subject Classification.* Primary: 03E35, 54D20. Secondary: 54C50, 03E05.

Key words and phrases. H -separable, M -separable, Laver forcing, Menger space, Hurewicz space.

Menger's property, as it is explicitly stated in the first paragraph of [10], was an application in dimension theory, one of the areas of interest of Mardešić. However, this paper concentrates on set-theoretic and combinatorial aspects of the property of Menger and its variations.

Theorem 1.1 is closely related to the main result of [13] asserting that in the Laver model the product of any two Hurewicz metrizable spaces has the Menger property. Let us note that our proof in [13] is conceptually different, even though both proofs are based on the same main technical lemma of [9]. Regarding the relation between Theorem 1.1 and the main result of [13], each of them implies a weak form of the other one via the following duality results: For a metrizable space X , $C_p(X)$ is M -separable (resp. H -separable) if and only if all finite powers of X are Menger (resp. Hurewicz), see [14, Theorem 35] and [3, Theorem 40], respectively. Thus Theorem 1.1 (combined with the well-known fact that $C_p(X)$ is hereditarily separable for metrizable separable spaces X) implies that in the Laver model, if all finite powers of metrizable separable spaces X_0, X_1 are Hurewicz, then $X_0 \times X_1$ is Menger. And vice versa: The main result of [13] implies that in the Laver model, the product of two H -separable spaces of the form $C_p(X)$ for a metrizable separable X , is M -separable.

The proof of Theorem 1.1, which is based on the analysis of names for reals in the style of [9], unfortunately seems to be rather tailored for the H -separability and we were not able to prove any analogous results even for small variations thereof. Recall from [6] that a space X is said to be wH -separable if for any *decreasing* sequence $\langle D_n : n \in \omega \rangle$ of dense subsets of X , one can pick finite subsets $F_n \subset D_n$ such that for any non-empty open $U \subset X$ the set $\{n \in \omega : U \cap F_n \neq \emptyset\}$ is co-finite. It is clear that every H -separable space is wH -separable, and it seems to be unknown whether the converse is (at least consistently) true. Combining [6, Lemma 2.7(2) and Corollary 4.2] we obtain that every countable Fréchet-Urysohn space is wH -separable, and to our best knowledge it is open whether countable Fréchet-Urysohn spaces must be H -separable. The statement “finite products of countable Fréchet-Urysohn spaces are M -separable” is known to be independent from ZFC: It follows from the PFA by [2, Theorem 3.3], holds in the Cohen model by [2, Corollary 3.2], and fails under CH by [1, Theorem 2.24]. Moreover¹, CH implies the existence of two countable Fréchet-Urysohn H -separable topological groups whose product is not M -separable, see [11, Corollary 6.2]. These results motivate the following

- Question 1.2.** (1) Is it consistent that the product of two countable wH -separable spaces is M -separable? Does this statement hold in the Laver model?
- (2) Is the product of two countable Fréchet-Urysohn space M -separable in the Laver model?

¹We do not know whether the spaces constructed in the proof of [1, Theorem 2.24] are H -separable.

- (3) Is the product of three (finitely many) countable H -separable spaces M -separable in the Laver model?
- (4) Is the product of finitely many countable H -separable spaces H -separable in the Laver model?

2. PROOF OF THEOREM 1.1

We need the following

Definition 2.1. A topological space $\langle X, \tau \rangle$ is called *box-separable* if for every function R assigning to each countable family $\mathcal{U}$ of non-empty open subsets of X a sequence $R(\mathcal{U}) = \langle F_n : n \in \omega \rangle$ of finite non-empty subsets of X such that $\{n : F_n \subset U\}$ is infinite for every $U \in \mathcal{U}$, there exists $\mathcal{U} \subset [\tau \setminus \{\emptyset\}]^\omega$ of size $|\mathcal{U}| = \omega_1$ such that for all $U \in \tau \setminus \{\emptyset\}$ there exists $\mathcal{U} \in \mathcal{U}$ such that $\{n : R(\mathcal{U})(n) \subset U\}$ is infinite.

Any countable space is obviously box-separable under CH, which makes the latter notion uninteresting when considered in arbitrary ZFC models. However, as we shall see in Lemma 2.3, the box-separability becomes useful under $\mathfrak{b} > \omega_1$. Here $\mathfrak{b}$ denotes the minimal cardinality of a subspace X of ω^ω which is not eventually dominated by a single function, see [4] for more information on $\mathfrak{b}$ and other cardinal characteristics of the reals.

The following lemma is the key part of the proof of Theorem 1.1. We will use the notation from [9] with the only difference being that smaller conditions in a forcing poset are supposed to carry more information about the generic filter, and the ground model is denoted by V .

A subset C of ω_2 is called an ω_1 -club if it is unbounded and for every $\alpha \in \omega_2$ of cofinality ω_1 , if $C \cap \alpha$ is cofinal in α then $\alpha \in C$.

Lemma 2.2. *In the Laver model every countable H -separable space is box-separable.*

Proof. We work in $V[G_{\omega_2}]$, where G_{ω_2} is $\mathbb{P}_{\omega_2}$ -generic and $\mathbb{P}_{\omega_2}$ is the iteration of length ω_2 with countable supports of the Laver forcing, see [9] for details. Let us fix an H -separable space of the form $\langle \omega, \tau \rangle$ and a function R such as in the definition of box-separability. By a standard argument (see, e.g., the proof of [5, Lemma 5.10]) there exists an ω_1 -club $C \subset \omega_2$ such that for every $\alpha \in C$ the following conditions hold:

- (i) $\tau \cap V[G_\alpha] \in V[G_\alpha]$ and for every sequence $\langle D_n : n \in \omega \rangle \in V[G_\alpha]$ of dense subsets of $\langle \omega, \tau \rangle$ there exists a sequence $\langle K_n : n \in \omega \rangle \in V[G_\alpha]$ such that $K_n \in [D_n]^{<\omega}$ and for every $U \in \tau \setminus \emptyset$ the intersection $U \cap K_n$ is non-empty for all but finitely many $n \in \omega$;
- (ii) $R(\mathcal{U}) \in V[G_\alpha]$ for any $\mathcal{U} \in [\tau \setminus \{\emptyset\}]^\omega \cap V[G_\alpha]$; and
- (iii) For every $A \in \mathcal{P}(\omega) \cap V[G_\alpha]$ the interior $\text{Int}(A)$ also belongs to $V[G_\alpha]$.

By [9, Lemma 11] there is no loss of generality in assuming that $0 \in C$. We claim that $\mathcal{U} := [\tau \setminus \{\emptyset\}]^\omega \cap V$ is a witness for $\langle \omega, \tau \rangle$ being box-separable.

Suppose, contrary to our claim, that there exists $A \in \tau \setminus \{\emptyset\}$ such that $R(\mathcal{U})(n) \not\subset A$ for all but finitely many $n \in \omega$ and $\mathcal{U} \in \mathcal{U}$. Let $\dot{A}$ be a $\mathbb{P}_{\omega_2}$ -name for A and $p \in \mathbb{P}_{\omega_2}$ a condition forcing the above statement. Applying [9, Lemma 14] to the sequence $\langle \dot{a}_i : i \in \omega \rangle$ such that $\dot{a}_i = \dot{A}$ for all $i \in \omega$, we get a condition $p' \leq p$ such that $p'(0) \leq^0 p(0)$, and a finite set $\mathcal{U}_s \subset \mathcal{P}(\omega)$ for every $s \in p'(0)$ with $p'(0) \langle 0 \rangle \leq s$, such that for each $n \in \omega$, $s \in p'(0)$ with $p'(0) \langle 0 \rangle \leq s$, and for all but finitely many immediate successors t of s in $p'(0)$ we have

$$p'(0)_t \wedge p' \restriction [1, \omega_2) \Vdash \exists U \in \mathcal{U}_s (\dot{A} \cap n = U \cap n).$$

Of course, any $p'' \leq p'$ also has the property above with the same $\mathcal{U}_s$'s. However, the stronger p'' is, the more elements of $\mathcal{U}_s$ might play no role any more. Therefore throughout the rest of the proof we shall call $U \in \mathcal{U}_s$ *void* for $p'' \leq p'$ and $s \in p''(0)$, where $p''(0) \langle 0 \rangle \leq s$, if there exists $n \in \omega$ such that for all but finitely many immediate successors t of s in $p''(0)$ there is no $q \leq p''(0)_t \wedge p'' \restriction [1, \omega_2)$ with the property $q \Vdash \dot{A} \cap n = U \cap n$. Note that for any $p'' \leq p'$ and $s \in p''(0)$, $p''(0) \langle 0 \rangle \leq s$, there exists $U \in \mathcal{U}_s$ which is non-void for p'', s . Two cases are possible.

a) For every $p'' \leq p'$ there exists $s \in p''(0)$, $p''(0) \langle 0 \rangle \leq s$, and a non-void $U \in \mathcal{U}_s$ for p'', s such that $\text{Int}(U) \neq \emptyset$. In this case let $\mathcal{U} \in \mathcal{U}$ be any countable family containing $\{\text{Int}(U) : U \in \bigcup_{s \in p''(0), p''(0) \langle 0 \rangle \leq s} \mathcal{U}_s\} \setminus \{\emptyset\}$. It follows from the above that p forces $R(\mathcal{U})(k) \not\subset \dot{A}$ for all but finitely many $k \in \omega$. Let $p'' \leq p'$ and $m \in \omega$ be such that p'' forces $R(\mathcal{U})(k) \not\subset \dot{A}$ for all $k \geq m$. Fix a non-void U for p'', s , where $s \in p''(0)$ and $p''(0) \langle 0 \rangle \leq s$, such that $\text{Int}(U) \neq \emptyset$ (and hence $\text{Int}(U) \in \mathcal{U}$). It follows from the above that there exists $k \geq m$ such that $R(\mathcal{U})(k) \subset \text{Int}(U) \subset U$. Let $n \in \omega$ be such that $R(\mathcal{U})(k) \subset n$. By the definition of being non-void there are infinitely many immediate successors t of s in $p''(0)$ for which there exists $q_t \leq p''(0)_t \wedge p'' \restriction [1, \omega_2)$ with the property $q_t \Vdash \dot{A} \cap n = U \cap n$. Then for any q_t as above we have that q_t forces $R(\mathcal{U})(k) \subset \dot{A}$ because $R(\mathcal{U})(k) \subset U \cap n$, which contradicts the fact that $q_t \leq p''$ and $p'' \Vdash R(\mathcal{U})(k) \not\subset \dot{A}$.

b) There exists $p'' \leq p'$ such that for all $s \in p''(0)$, $p''(0) \langle 0 \rangle \leq s$, every $U \in \mathcal{U}_s$ with $\text{Int}(U) \neq \emptyset$ is void for p'', s . Note that this implies that every $U \in \mathcal{U}_s$ with $\text{Int}(U) \neq \emptyset$ is void for q, s for all $q \leq p''$ and $s \in q(0)$ such that $q(0) \langle 0 \rangle \leq s$.

Let $\langle D_k : k \in \omega \rangle \in V$ be a sequence of dense subsets of $\langle \omega, \tau \rangle$ such that for every $U \in \bigcup_{s \in p''(0), p''(0) \langle 0 \rangle \leq s} \mathcal{U}_s$, if $\text{Int}(U) = \emptyset$, then $\omega \setminus U = D_k$ for infinitely many $k \in \omega$. Let $\langle K_k : k \in \omega \rangle \in V$ be such as in item (i) above. Then p'' forces that $K_k \cap \dot{A} \neq \emptyset$ for all but finitely many $k \in \omega$. Passing to a stronger condition, we may additionally assume if necessary, that there exists $m \in \omega$ such that $p'' \Vdash \forall k \geq m (K_k \cap \dot{A} \neq \emptyset)$.

Fix $U \in \mathcal{U}_{p''(0) \langle 0 \rangle}$ non-void for $p'', p''(0) \langle 0 \rangle$. Then $\text{Int}(U) = \emptyset$ by the choice of p'' and hence there exists $k \geq m$ such that $\omega \setminus U = D_k$. It follows that $K_k \cap U = \emptyset$ because $K_k \subset D_k$. On the other hand, since U is non-void

for $p'', p''(0)\langle 0 \rangle$, for $n = \max K_k + 1$ we can find infinitely many immediate successors t of $p''(0)\langle 0 \rangle$ in $p''(0)$ for which there exists $q_t \leq p''(0)_t \hat{\wedge} p'' \upharpoonright [1, \omega_2)$ forcing $\dot{A} \cap n = U \cap n$. Then any such q_t forces $K_k \cap \dot{A} = \emptyset$ (because $K_k \subset n$ and $K_k \cap U = \emptyset$), contradicting the fact that $p'' \geq q_t$ and $p'' \Vdash K_k \cap \dot{A} \neq \emptyset$.

Contradictions obtained in cases *a*) and *b*) above imply that $\mathbf{U} := [\tau \setminus \{\emptyset\}]^\omega \cap V$ is a witness for $\langle \omega, \tau \rangle$ being box-separable, which completes our proof. $\square$

Theorem 1.1 is a direct consequence of Lemma 2.2 combined with the following

Lemma 2.3. *Suppose that $\mathfrak{b} > \omega_1$, X is box-separable, and Y is H -separable. Then $X \times Y$ is M -separable provided that it is separable.*

Proof. Let $\langle D_n : n \in \omega \rangle$ be a sequence of countable dense subsets of $X \times Y$. Let us fix a countable family $\mathcal{U}$ of open non-empty subsets of X and a partition $\omega = \sqcup_{U \in \mathcal{U}} \Omega_U$ into infinite pieces. For every $n \in \Omega_U$ set $D_n^\mathcal{U} = \{y \in Y : \exists x \in U (\langle x, y \rangle \in D_n)\}$ and note that $D_n^\mathcal{U}$ is dense in Y for all $n \in \omega$. Therefore there exists a sequence $\langle L_n^\mathcal{U} : n \in \omega \rangle$ such that $L_n^\mathcal{U} \in [D_n^\mathcal{U}]^{<\omega}$ and for every open non-empty $V \subset Y$ we have $L_n^\mathcal{U} \cap V \neq \emptyset$ for all but finitely many n . For every $n \in \Omega_U$ find $K_n^\mathcal{U} \in [U]^{<\omega}$ such that for every $y \in L_n^\mathcal{U}$ there exists $x \in K_n^\mathcal{U}$ such that $\langle x, y \rangle \in D_n$, and set $R(\mathcal{U}) = \langle K_n^\mathcal{U} : n \in \omega \rangle$. Note that R is such as in the definition of box-separability because $K_n^\mathcal{U} \subset U$ for all $n \in \Omega_U$ and the latter set is infinite. Since X is box-separable there exists a family $\mathbf{U}$ of countable collections of open non-empty subsets of X of size $|\mathbf{U}| = \omega_1$, and such that for every open non-empty $U \subset X$ there exists $\mathcal{U} \in \mathbf{U}$ with the property $R(\mathcal{U})(n) \subset U$ for infinitely many n . Since each D_n is countable and $|\mathbf{U}| < \mathfrak{b}$, there exists a sequence $\langle F_n : n \in \omega \rangle$ such that $F_n \in [D_n]^{<\omega}$ and for every $\mathcal{U} \in \mathbf{U}$ we have $F_n \supset (K_n^\mathcal{U} \times L_n^\mathcal{U}) \cap D_n$ for all but finitely many $n \in \omega$.

We claim that $\bigcup_{n \in \omega} F_n$ is dense in $X \times Y$. Indeed, let us fix open non-empty subset of $X \times Y$ of the form $U \times V$ and find $\mathcal{U} \in \mathbf{U}$ with the property $R(\mathcal{U})(n) = K_n^\mathcal{U} \subset U$ for infinitely many n , say for all $n \in I \in [\omega]^\omega$. Passing to a co-finite subset of I , we may assume if necessary, that $F_n \supset (K_n^\mathcal{U} \times L_n^\mathcal{U}) \cap D_n$ for all $n \in I$. Finally, fix $n \in I$ such that $L_n^\mathcal{U} \cap V \neq \emptyset$ and pick $y \in L_n^\mathcal{U} \cap V$. By the definition of $D_n^\mathcal{U}$ and $L_n^\mathcal{U} \subset D_n^\mathcal{U}$ we can find $x \in K_n^\mathcal{U}$ such that $\langle x, y \rangle \in D_n$. Then $\langle x, y \rangle \in U \times V$ and $\langle x, y \rangle \in F_n$ because $\langle x, y \rangle \in K_n^\mathcal{U} \times L_n^\mathcal{U}$ and $\langle x, y \rangle \in D_n$. This completes our proof. $\square$

ACKNOWLEDGEMENTS

This research was supported by the Slovenian Research Agency grants P1-0292, J1-8131, J1-7025, J1-6721, and J1-5435, and the Austrian Science Fund FWF (Grants I 1209-N25 and I 2374-N35).

REFERENCES

- [1] Barman, D.; Dow, A., *Selective separability and $SS+$* , Topology Proc. **37** (2011), 181–204.
- [2] Barman, D.; Dow, A., *Proper forcing axiom and selective separability*, Topology Appl. **159** (2012), 806–813.
- [3] Bella, A., Bonanzinga, M., Matveev, M., *Variations of selective separability*, Topology Appl. **156** (2009), 1241–1252.
- [4] Blass, A., *Combinatorial cardinal characteristics of the continuum*, in: *Handbook of Set Theory* (M. Foreman, A. Kanamori, and M. Magidor, eds.), Springer, 2010, pp. 395–491.
- [5] Blass, A.; Shelah, S., *There may be simple $P_{\aleph_1}$ - and $P_{\aleph_2}$ -points and the Rudin-Keisler ordering may be downward directed*, Ann. Pure Appl. Logic **33** (1987), 213–243.
- [6] Gruenhage, G.; Sakai, M., *Selective separability and its variations*, Topology Appl. **158** (2011) 1352–1359.
- [7] Hurewicz, W., *Über die Verallgemeinerung des Borellschen Theorems*, Math. Z. **24** (1925), 401–421.
- [8] Hurewicz, W., *Über Folgen stetiger Funktionen*, Fund. Math. **9** (1927), 193–204.
- [9] Laver, R., *On the consistency of Borel’s conjecture*, Acta Math. **137** (1976), 151–169.
- [10] Menger, K., *Einige Überdeckungssätze der Punktmengenlehre*, Sitzungsberichte. Abt. 2a, Mathematik, Astronomie, Physik, Meteorologie und Mechanik (Wiener Akademie) **133** (1924), 421–444.
- [11] Miller, A.W.; Tsaban, B.; Zdomskyy, L., *Selective covering properties of product spaces, II: gamma spaces*, Trans. Amer. Math. Soc. **368** (2016), 2865–2889.
- [12] Repovš, D.; Zdomskyy, L., *On M -separability of countable spaces and function spaces*, Topology Appl. **157** (2010), 2538–2541.
- [13] Repovš, D.; Zdomskyy, L., *Products of Hurewicz spaces in the Laver model*, Bull. Symb. Log. **23:3** (2017), 324–333.
- [14] Scheepers, M., *Combinatorics of open covers. VI. Selectors for sequences of dense sets*, Quaest. Math. **22** (1999), 109–130.

FACULTY OF EDUCATION AND MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA & INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, 1000 LJUBLJANA, SLOVENIA.

E-mail address: dusan.repovs@guest.arnes.si

URL: <http://www.fmf.uni-lj.si/~repovs/index.htm>

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, UNIVERSITY OF VIENNA, A-1090 WIEN, AUSTRIA.

E-mail address: lzdomsky@gmail.com

URL: <http://www.logic.univie.ac.at/~lzdomsky/>