

ON MONOIDS OF INJECTIVE PARTIAL COFINITE SELFMAPS

OLEG GUTIK AND DUŠAN REPOVŠ

ABSTRACT. We study the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ of injective partial cofinite selfmaps of an infinite cardinal λ . We show that $\mathcal{J}_\lambda^{\text{cf}}$ is a bisimple inverse semigroup and each chain of idempotents in $\mathcal{J}_\lambda^{\text{cf}}$ is contained in a bicyclic subsemigroup of $\mathcal{J}_\lambda^{\text{cf}}$, we describe the Green relations on $\mathcal{J}_\lambda^{\text{cf}}$ and we prove that every non-trivial congruence on $\mathcal{J}_\lambda^{\text{cf}}$ is a group congruence. Also, we describe the structure of the quotient semigroup $\mathcal{J}_\lambda^{\text{cf}}/\sigma$, where σ is the least group congruence on $\mathcal{J}_\lambda^{\text{cf}}$.

1. INTRODUCTION AND PRELIMINARIES

In this paper we shall denote the first infinite ordinal by ω and the cardinality of the set A by $|A|$. We shall identify all cardinals with their corresponding initial ordinals. We shall denote the set of integers by \mathbb{Z} and the additive group of integers by $\mathbb{Z}(+)$.

A semigroup S is called *inverse* if for every element $x \in S$ there exists a unique $x^{-1} \in S$ such that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. The element x^{-1} is called the *inverse of $x \in S$* . If S is an inverse semigroup, then the function $\text{inv}: S \rightarrow S$ which assigns to every element x of S its inverse element x^{-1} is called the *inversion*.

A congruence \mathfrak{C} on a semigroup S is called *non-trivial* if \mathfrak{C} is distinct from the universal and the identity congruences on S , and a *group congruence* if the quotient semigroup S/\mathfrak{C} is a group.

If S is a semigroup, then we shall denote the subset of all idempotents in S by $E(S)$. If S is an inverse semigroup, then $E(S)$ is closed under multiplication and we shall refer to $E(S)$ as a *band* (or the *band of S*). Then the semigroup operation on S determines the following partial order \leq on $E(S)$: $e \leq f$ if and only if $ef = fe = e$. This order is called the *natural partial order* on $E(S)$. A *semilattice* is a commutative semigroup of idempotents. A semilattice E is called *linearly ordered* or a *chain* if its natural order is a linear order. A *maximal chain* of a semilattice E is a chain which is not properly contained in any other chain of E .

The Axiom of Choice implies the existence of maximal chains in every partially ordered set. According to [13, Definition II.5.12], a chain L is called an ω -*chain* if L is isomorphic to $\{0, -1, -2, -3, \dots\}$ with the usual order \leq or equivalently, if L is isomorphic to (ω, \max) . Let E be a semilattice and $e \in E$. We put $\downarrow e = \{f \in E \mid f \leq e\}$ and $\uparrow e = \{f \in E \mid e \leq f\}$. By $(\mathcal{P}_{<\omega}(\lambda), \cup)$ we shall denote the *free semilattice with identity* over a set of cardinality $\lambda \geq \omega$, i.e., $(\mathcal{P}_{<\omega}(\lambda), \cup)$ is the set of all finite subsets (with the empty set) of λ with the semilattice operation “union”.

2010 *Mathematics Subject Classification*. Primary 20M20, 20M18; Secondary 20B30.

Key words and phrases. Bicyclic semigroup, semigroup of bijective partial transformations, congruence, symmetric group, group congruence, semidirect product.

If S is a semigroup, then we shall denote the Green relations on S by \mathcal{R} , \mathcal{L} , \mathcal{J} , \mathcal{D} and \mathcal{H} (see [5]). A semigroup S is called *simple* if S does not contain proper two-sided ideals and *bisimple* if S has only one \mathcal{D} -class.

The bicyclic semigroup $\mathcal{C}(p, q)$ is the semigroup with the identity 1 generated by elements p and q subject only to the condition $pq = 1$. The bicyclic semigroup is bisimple and every one of its congruences is either trivial or a group congruence. Moreover, every homomorphism h of the bicyclic semigroup is either an isomorphism or the image of $\mathcal{C}(p, q)$ under h is a cyclic group (see [5, Corollary 1.32]). The bicyclic semigroup plays an important role in algebraic theory of semigroups and in the theory of topological semigroups. For example a well-known Andersen's result [1] states that a (0-)simple semigroup is completely (0-)simple if and only if it does not contain the bicyclic semigroup. The bicyclic semigroup admits only the discrete topology [7]. The problem of embeddability of the bicycle semigroup into compact-like semigroups was studied in [2, 3, 4, 8, 11].

Remark 1.1. *We observe that the bicyclic semigroup is isomorphic to the semigroup $\mathcal{C}_{\mathbb{N}}(\alpha, \beta)$ which is generated by partial transformations α and β of the set of positive integers \mathbb{N} , defined as follows: $(n)\alpha = n + 1$ if $n \geq 1$ and $(n)\beta = n - 1$ if $n > 1$ (see Exercise IV.1.11(ii) in [13]).*

If T is a semigroup, then we say that a subsemigroup S of T is a *bicyclic subsemigroup* of T if S is isomorphic to the bicyclic semigroup $\mathcal{C}(p, q)$.

Hereafter we shall assume that λ is an infinite cardinal. If $\alpha: X \rightarrow Y$ is a partial map, then we shall denote the domain and the range of α by $\text{dom } \alpha$ and $\text{ran } \alpha$, respectively.

Let \mathcal{J}_λ denote the set of all partial one-to-one transformations of an infinite set X of cardinality λ together with the following semigroup operation:

$$x(\alpha\beta) = (x\alpha)\beta \quad \text{if } x \in \text{dom}(\alpha\beta) = \{y \in \text{dom } \alpha \mid y\alpha \in \text{dom } \beta\}, \quad \text{for } \alpha, \beta \in \mathcal{J}_\lambda.$$

The semigroup \mathcal{J}_λ is called the *symmetric inverse semigroup* over the set X (see [5, Section 1.9]). The symmetric inverse semigroup was introduced by Vagner [21] and it plays a major role in the theory of semigroups.

Furthermore, we shall identify the cardinal $\lambda = |X|$ with the set X . By $\mathcal{J}_\lambda^{\text{cf}}$ we shall denote a subsemigroup of injective partial selfmaps of λ with cofinite domains and ranges in \mathcal{J}_λ , i.e.,

$$\mathcal{J}_\lambda^{\text{cf}} = \{\alpha \in \mathcal{J}_\lambda \mid |\lambda \setminus \text{dom } \alpha| < \infty \quad \text{and} \quad |\lambda \setminus \text{ran } \alpha| < \infty\}.$$

Obviously, $\mathcal{J}_\lambda^{\text{cf}}$ is an inverse submonoid of the semigroup \mathcal{J}_λ . We shall call the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ the *monoid of injective partial cofinite selfmaps* of λ .

Next, by \mathbb{I} we shall denote the identity and by $H(\mathbb{I})$ the group of units of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$.

It well known that each partial injective cofinite selfmap f of λ induces a homeomorphism $f^*: \lambda^* \rightarrow \lambda^*$ of the remainder $\lambda^* = \beta\lambda \setminus \lambda$ of the Stone-Ćech compactification of the discrete space λ . Moreover, under some set theoretic axioms (like **PFA** or **OCA**), each homeomorphism of ω^* is induced by some partial injective cofinite selfmap of ω (see [15]–[20]). So the inverse semigroup $\mathcal{J}_\lambda^{\text{cf}}$ admits a natural homomorphism $\mathfrak{h}: \mathcal{J}_\lambda^{\text{cf}} \rightarrow \mathcal{H}(\lambda^*)$ to the homeomorphism group $\mathcal{H}(\lambda^*)$ of λ^* and this homomorphism is surjective under certain set theoretic assumptions.

The semigroups $\mathcal{J}_\infty^{\text{cf}}(\mathbb{N})$ and $\mathcal{J}_\infty^{\text{cf}}(\mathbb{Z})$ of injective isotone partial selfmaps with cofinite domains and images of positive integers and integers, respectively, were studied

in [9] and [10]. There it was proved that the semigroups $\mathcal{J}_\infty^\rightarrow(\mathbb{N})$ and $\mathcal{J}_\infty^\rightarrow(\mathbb{Z})$ have properties similar to the bicyclic semigroup: they are bisimple and every non-trivial homomorphic image of $\mathcal{J}_\infty^\rightarrow(\mathbb{N})$ and $\mathcal{J}_\infty^\rightarrow(\mathbb{Z})$ is a group, and moreover, the semigroup $\mathcal{J}_\infty^\rightarrow(\mathbb{N})$ has $\mathbb{Z}(+)$ as a maximal group image and $\mathcal{J}_\infty^\rightarrow(\mathbb{Z})$ has $\mathbb{Z}(+) \times \mathbb{Z}(+)$, respectively.

In this paper we shall study algebraic properties of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. We shall show that $\mathcal{J}_\lambda^{\text{cf}}$ is a bisimple inverse semigroup and every chain of idempotents in $\mathcal{J}_\lambda^{\text{cf}}$ is contained in a bicyclic subsemigroup of $\mathcal{J}_\lambda^{\text{cf}}$, we shall describe the Green relations on $\mathcal{J}_\lambda^{\text{cf}}$ and we shall prove that every non-trivial congruence on $\mathcal{J}_\lambda^{\text{cf}}$ is a group congruence. Also, we shall describe the structure of the quotient semigroup $\mathcal{J}_\lambda^{\text{cf}}/\sigma$, where σ is the least group congruence on $\mathcal{J}_\lambda^{\text{cf}}$.

2. ALGEBRAIC PROPERTIES OF THE SEMIGROUP $\mathcal{J}_\lambda^{\text{cf}}$

Proposition 2.1. (i) $\mathcal{J}_\lambda^{\text{cf}}$ is a simple semigroup.

- (ii) An element α of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is an idempotent if and only if $(x)\alpha = x$ for every $x \in \text{dom } \alpha$.
- (iii) If $\varepsilon, \iota \in E(\mathcal{J}_\lambda^{\text{cf}})$, then $\varepsilon \leq \iota$ if and only if $\text{dom } \varepsilon \subseteq \text{dom } \iota$.
- (iv) The semilattice $E(\mathcal{J}_\lambda^{\text{cf}})$ is isomorphic to $(\mathcal{P}_{<\omega}(\lambda), \cup)$ under the mapping $(\varepsilon)h = \lambda \setminus \text{dom } \varepsilon$.
- (v) Every maximal chain in $E(\mathcal{J}_\lambda^{\text{cf}})$ is an ω -chain.
- (vi) $\alpha\mathcal{R}\beta$ in $\mathcal{J}_\lambda^{\text{cf}}$ if and only if $\text{dom } \alpha = \text{dom } \beta$.
- (vii) $\alpha\mathcal{L}\beta$ in $\mathcal{J}_\lambda^{\text{cf}}$ if and only if $\text{ran } \alpha = \text{ran } \beta$.
- (viii) $\alpha\mathcal{H}\beta$ in $\mathcal{J}_\lambda^{\text{cf}}$ if and only if $\text{dom } \alpha = \text{dom } \beta$ and $\text{ran } \alpha = \text{ran } \beta$.
- (ix) $\alpha\mathcal{D}\beta$ for all $\alpha, \beta \in \mathcal{J}_\lambda^{\text{cf}}$ and hence the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is bisimple.

Proof. (i) We shall show that $\mathcal{J}_\lambda^{\text{cf}} \cdot \alpha \cdot \mathcal{J}_\lambda^{\text{cf}} = \mathcal{J}_\lambda^{\text{cf}}$ for every element $\alpha \in \mathcal{J}_\lambda^{\text{cf}}$. Let α and β be arbitrary elements of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. We shall choose elements $\gamma, \delta \in \mathcal{J}_\lambda^{\text{cf}}$ such that $\gamma \cdot \alpha \cdot \delta = \beta$. We put $\text{dom } \gamma = \text{dom } \beta$, $\text{ran } \gamma = \text{dom } \alpha$, $\text{dom } \delta = \text{ran } \alpha$ and $\text{ran } \delta = \text{ran } \beta$. Since the sets $\lambda \setminus \text{dom } \alpha$ and $\lambda \setminus \text{dom } \beta$ are finite we conclude that there exists a bijective map $f: \text{dom } \alpha \rightarrow \text{dom } \beta$. We put $\gamma = f$ and $((x)\gamma)\alpha\delta = (x)\beta$ for all $x \in \text{dom } \beta$. Then we have that $\gamma \cdot \alpha \cdot \delta = \beta$.

Statements (ii) – (v) are trivial and they follow from the definition of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. The proofs of (vi) – (viii) follow trivially from the fact that $\mathcal{J}_\lambda^{\text{cf}}$ is a regular semigroup, and Proposition 2.4.2 and Exercise 5.11.2 in [12].

(ix) Let α and β be arbitrary elements of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. Since the sets $\lambda \setminus \text{dom } \alpha$ and $\lambda \setminus \text{ran } \beta$ are finite we conclude that there exists a bijective map $\gamma: \text{dom } \alpha \rightarrow \text{ran } \beta$. Then $\gamma \in \mathcal{J}_\lambda^{\text{cf}}$ and by statements (vi) and (vii) we have that $\alpha\mathcal{R}\gamma$ and $\beta\mathcal{L}\gamma$ in $\mathcal{J}_\lambda^{\text{cf}}$ and hence $\alpha\mathcal{D}\beta$ in $\mathcal{J}_\lambda^{\text{cf}}$. \square

We denote the group of all bijective transformations of a set of cardinality λ by \mathcal{S}_λ . Then we get the following:

Corollary 2.2. The group of units $H(\mathbb{I})$ of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is isomorphic to \mathcal{S}_λ .

For any idempotents ε and ι of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ we denote:

$$H(\varepsilon, \iota) = \{\chi \in \mathcal{J}_\lambda^{\text{cf}} \mid \chi \cdot \chi^{-1} = \varepsilon \text{ and } \chi^{-1} \cdot \chi = \iota\} \quad \text{and} \quad H(\varepsilon) = H(\varepsilon, \varepsilon).$$

Proposition 2.1(viii) implies that the set $H(\varepsilon, \iota)$ is a \mathcal{H} -class and the set $H(\varepsilon)$ is a maximal subgroup in $\mathcal{J}_\lambda^{\text{cf}}$ for all idempotents $\varepsilon, \iota \in \mathcal{J}_\lambda^{\text{cf}}$.

Corollary 2.2 and Proposition 2.20 of [5] imply the following:

Corollary 2.3. *Every maximal subgroup of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is isomorphic to \mathcal{S}_λ .*

Proposition 2.4. $|\mathcal{J}_\lambda^{\text{cf}}| = 2^{|\lambda|}$.

Proof. Since $|\lambda \times \lambda| = |\lambda|$ we have that $|\mathcal{S}_\lambda| \leq 2^{|\lambda \times \lambda|} = 2^{|\lambda|}$. Since $|\lambda \sqcup \lambda| = |\lambda|$ there exists an injective map $f: \mathcal{P}(\lambda) \rightarrow \mathcal{S}_{\lambda \sqcup \lambda}$ from the set $\mathcal{P}(\lambda)$ of all subset of the cardinal λ into the group $\mathcal{S}_{\lambda \sqcup \lambda}$ defined in the following way: $f(A)$ is a bijection on $\lambda \sqcup \lambda$ with support $A \sqcup A$. Then we have that $|\mathcal{S}_\lambda| \geq 2^{|\lambda \sqcup \lambda|} = 2^{|\lambda|}$ and hence $|\mathcal{S}_\lambda| = 2^{|\lambda|}$.

Since $|\mathcal{P}_{<\omega}(\lambda)| = |\mathcal{P}_{<\omega}(\lambda) \times \mathcal{P}_{<\omega}(\lambda)| = \lambda$ we conclude that Theorem 2.20 from [5] and Proposition 2.1(viii) imply that

$$|\mathcal{J}_\lambda^{\text{cf}}| = |\mathcal{P}_{<\omega}(\lambda) \times \mathcal{P}_{<\omega}(\lambda) \times \mathcal{S}_\lambda| = |\mathcal{P}_{<\omega}(\lambda) \times \mathcal{P}_{<\omega}(\lambda)| \cdot |\mathcal{S}_\lambda| = |\lambda| \cdot 2^{|\lambda|} = 2^{|\lambda|}.$$

□

Proposition 2.5. *For every $\alpha, \beta \in \mathcal{J}_\lambda^{\text{cf}}$, both sets*

$$\{\chi \in \mathcal{J}_\lambda^{\text{cf}} \mid \alpha \cdot \chi = \beta\} \quad \text{and} \quad \{\chi \in \mathcal{J}_\lambda^{\text{cf}} \mid \chi \cdot \alpha = \beta\}$$

are finite. Consequently, every right translation and every left translation by an element of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is a finite-to-one map.

Proof. We denote

$$A = \{\chi \in \mathcal{J}_\lambda^{\text{cf}} \mid \alpha \cdot \chi = \beta\} \quad \text{and} \quad B = \{\chi \in \mathcal{J}_\lambda^{\text{cf}} \mid \alpha^{-1} \cdot \alpha \cdot \chi = \alpha^{-1} \cdot \beta\}.$$

Then $A \subseteq B$ and the restriction of any partial map $\chi \in B$ onto $\text{dom}(\alpha^{-1} \cdot \alpha)$ coincides with the partial map $\alpha^{-1} \cdot \beta$. Since every partial map from $\mathcal{J}_\lambda^{\text{cf}}$ has cofinite range and cofinite domain we conclude that the set B is finite and hence so is A . □

Proposition 2.6. *Each maximal chain L of idempotents in $\mathcal{J}_\lambda^{\text{cf}}$ coincides with the idempotent band $E(S)$ of a bicyclic subsemigroup S of $\mathcal{J}_\lambda^{\text{cf}}$.*

Proof. By Proposition 2.1(iii), the chain L can be written as $L = \{\varepsilon_n\}_{n=1}^\infty$ where $\varepsilon_1 > \varepsilon_2 > \dots > \varepsilon_n > \dots$. Since every infinite subchain of an ω -chain is also an ω -chain we have that Proposition 2.1(v) implies that L is an ω -chain. Then by Proposition 2.1(iii) we get that $\text{dom } \varepsilon_i \setminus \text{dom } \varepsilon_{i+1} \neq \emptyset$ for all positive integers i . Also, the maximality of L implies that the set $\text{dom } \varepsilon_i \setminus \text{dom } \varepsilon_{i+1}$ is a singleton for all positive integers i . For every positive integer i we put $\{x_i\} = \text{dom } \varepsilon_i \setminus \text{dom } \varepsilon_{i+1}$. Then we put $D = \text{dom } \varepsilon_1 \setminus \bigcup_{i \in \mathbb{N}} \{x_i\}$ and define the partial maps $\alpha: \lambda \rightarrow \lambda$ and $\beta: \lambda \rightarrow \lambda$ as follows:

$$(x)\alpha = \begin{cases} x_{n+1}, & \text{if } x = x_n \in \text{dom } \varepsilon_1 \setminus D \text{ and } n \geq 1; \\ x, & \text{if } x \in D; \end{cases}$$

and

$$(x)\beta = \begin{cases} x_{n-1}, & \text{if } x = x_n \in \text{dom } \varepsilon_1 \setminus D \text{ and } n > 1; \\ x, & \text{if } x \in D. \end{cases}$$

Since the set $\lambda \setminus \text{dom } \varepsilon_1$ is finite we have that $\alpha, \beta \in \mathcal{J}_\lambda^{\text{cf}}$ and Remark 1.1 implies the statement of our proposition. □

Proposition 2.6 and the Axiom of Choice imply the following proposition.

Proposition 2.7. *Each chain of idempotents in $\mathcal{J}_\lambda^{\text{cf}}$ is contained in a bicyclic subsemigroup of $\mathcal{J}_\lambda^{\text{cf}}$.*

Proposition 2.8. *Let \mathfrak{C} be a congruence on the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. If there exist two non- \mathcal{H} -equivalent elements $\alpha, \beta \in \mathcal{J}_\lambda^{\text{cf}}$ such that $\alpha\mathfrak{C}\beta$, then \mathfrak{C} is a group congruence on $\mathcal{J}_\lambda^{\text{cf}}$.*

Proof. First we suppose that α and β are distinct idempotents of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. Without loss of generality we can assume that α and β are compatible and $\alpha \leq \beta$. Otherwise, replace α by $\alpha \cdot \beta$. Then by Proposition 2.7 there exists a maximal chain L in $E(\mathcal{J}_\lambda^{\text{cf}})$ such that L contains the elements α and β , and hence L contained in a bicyclic subsemigroup S of $\mathcal{J}_\lambda^{\text{cf}}$. Then Corollary 1.32 of [5] implies that $\varepsilon\mathfrak{C}\iota$ for all elements ε and ι of the chain L .

Let ν be an arbitrary idempotent of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. Obviously, if $\varepsilon, \iota \in L$ such that $\varepsilon \leq \iota$ then $\varepsilon \cdot \nu \leq \iota \cdot \nu$. Since $\uparrow e$ is a finite subset of the free semilattice with unity $(\mathcal{P}_{<\omega}(\lambda), \subseteq)$ for any $e \in (\mathcal{P}_{<\omega}(\lambda), \subseteq)$, we have that Proposition 2.1(iv) implies that νL is an infinite chain in $E(\mathcal{J}_\lambda^{\text{cf}})$. Then we have that $\varepsilon\mathfrak{C}\iota$ for all $\varepsilon, \iota \in \nu L$. We put $L_\nu = \nu L \cup \{\nu\} \cup \{\mathbb{I}\}$. Then L_ν is a chain in $E(\mathcal{J}_\lambda^{\text{cf}})$. Therefore by Proposition 2.7 we get that there exists a maximal chain L_{\max} in $E(\mathcal{J}_\lambda^{\text{cf}})$ which contains the chain L_ν and L_{\max} is a band of a bicyclic subsemigroup S in $\mathcal{J}_\lambda^{\text{cf}}$. Now Corollary 1.32 of [5] implies that $\varepsilon\mathfrak{C}\iota$ for all elements ε and ι of the chain L_ν . Hence $\nu\mathfrak{C}\mathbb{I}$ and $\alpha\mathfrak{C}\mathbb{I}$ imply that $\nu\mathfrak{C}\alpha$. Therefore all idempotents of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ are \mathfrak{C} -equivalent. Since the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is inverse we conclude that quotient semigroup $\mathcal{J}_\lambda^{\text{cf}}/\mathfrak{C}$ contains only one idempotent and by Lemma II.1.10 from [13] the semigroup $\mathcal{J}_\lambda^{\text{cf}}/\mathfrak{C}$ is a group.

Suppose that α and β are distinct non- \mathcal{H} -equivalent elements of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ such that $\alpha\mathfrak{C}\beta$. Then Proposition 2.1 implies that at least one of the following conditions holds:

$$\alpha\alpha^{-1} \neq \beta\beta^{-1} \quad \text{or} \quad \alpha^{-1}\alpha \neq \beta^{-1}\beta.$$

By Lemma III.1.1 from [13] we have that $\alpha^{-1}\mathfrak{C}\beta^{-1}$. Then $\alpha\alpha^{-1}\mathfrak{C}\alpha\beta^{-1}$ and $\beta\beta^{-1}\mathfrak{C}\alpha\beta^{-1}$ and hence $\alpha\alpha^{-1}\mathfrak{C}\beta\beta^{-1}$. Similarly we get that $\alpha^{-1}\alpha\mathfrak{C}\beta^{-1}\beta$. Then the first part of the proof implies that \mathfrak{C} is a group congruence on $\mathcal{J}_\lambda^{\text{cf}}$. \square

Theorem 2.9. *Every non-trivial congruence on the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ is a group congruence.*

Proof. Let \mathfrak{C} be a non-trivial congruence on the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. Let α and β be distinct \mathfrak{C} -equivalent elements of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$. If the elements α and β are not \mathcal{H} -equivalent then Proposition 2.8 implies the statement of the theorem.

Suppose that $\alpha\mathcal{H}\beta$. Then Theorem 2.20 from [5] implies that without loss of generality we can assume that α and β are elements of the group of units $H(\mathbb{I})$ of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ and hence $\mathbb{I}\mathfrak{C}(\beta\alpha^{-1})$. We denote $\gamma = \beta\alpha^{-1}$. Since $\mathbb{I} \neq \gamma$ we conclude that there exists $x_0 \in \lambda$ such that $(x_0)\gamma \neq x_0$. We define ε to be an identity selfmap of the set $\lambda \setminus \{x_0\}$. Then $\varepsilon \in \mathcal{J}_\lambda^{\text{cf}}$ and $(\varepsilon \cdot \mathbb{I})\mathfrak{C}(\varepsilon \cdot \gamma)$. Since $(x_0)\gamma \neq x_0$ we have that Proposition 2.1(viii) implies that the elements ε and $\varepsilon \cdot \gamma$ are not \mathcal{H} -equivalent. Then by Proposition 2.8 we get that \mathfrak{C} is a group congruence on $\mathcal{J}_\lambda^{\text{cf}}$. \square

3. ON THE LEAST GROUP CONGRUENCE ON THE SEMIGROUP $\mathcal{J}_\lambda^{\text{cf}}$

Every inverse semigroup S admits the least group congruence σ (see [13, Section III]):

$$s\sigma t \quad \text{if and only if} \quad \text{there exists an idempotent } e \in S \quad \text{such that} \quad se = te.$$

Theorem 2.9 implies that every non-injective homomorphism $h: \mathcal{J}_\lambda^{\text{cf}} \rightarrow S$ from the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ into an arbitrary semigroup S generates a group congruence \mathfrak{h} on $\mathcal{J}_\lambda^{\text{cf}}$. In this section we describe the structure of the quotient semigroup $\mathcal{J}_\lambda^{\text{cf}}/\sigma$.

Proposition 3.1. *If $\alpha\sigma\beta$ in $\mathcal{J}_\lambda^{\text{cf}}$ then*

$$|\lambda \setminus \text{dom } \alpha| - |\lambda \setminus \text{ran } \alpha| = |\lambda \setminus \text{dom } \beta| - |\lambda \setminus \text{ran } \beta|.$$

Proof. Let ε be an idempotent of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ such that $\alpha\varepsilon = \beta\varepsilon$. We shall show that the statement of the proposition holds for the elements α and $\alpha\varepsilon$.

Without loss of generality we can assume that $\varepsilon \leq \alpha^{-1}\alpha$, i.e., $\text{dom } \varepsilon \subseteq \text{dom}(\alpha^{-1}\alpha)$. Since α is an injective partial map with $|\lambda \setminus \text{dom } \alpha| < \infty$ and $|\lambda \setminus \text{ran } \alpha| < \infty$, and ε is an identity map of the cofinite subset $\text{dom } \varepsilon$ in λ we conclude that

$$|\lambda \setminus \text{dom } \alpha| - |\lambda \setminus \text{ran } \alpha| = |\lambda \setminus \text{dom}(\alpha\varepsilon)| - |\lambda \setminus \text{ran}(\alpha\varepsilon)|.$$

This implies the statement of the proposition. \square

For an arbitrary element α of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ we denote

$$\bar{d}(\alpha) = |\lambda \setminus \text{dom } \alpha| \quad \text{and} \quad \bar{r}(\alpha) = |\lambda \setminus \text{ran } \alpha|.$$

Proposition 3.2. *If α and β are arbitrary elements of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ then*

$$\bar{d}(\alpha\beta) - \bar{r}(\alpha\beta) = \bar{d}(\alpha) - \bar{r}(\alpha) + \bar{d}(\beta) - \bar{r}(\beta).$$

Proof. We consider four cases.

(1) First we consider the case when $\text{ran } \alpha \subseteq \text{dom } \beta$. We put $k = \bar{r}(\alpha) - \bar{d}(\beta)$. Then the definition of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ implies that $k \geq 0$, $\bar{d}(\alpha\beta) = \bar{d}(\alpha)$, $\bar{r}(\alpha\beta) = \bar{r}(\beta) - k$, and hence in this case we get that

$$\bar{d}(\alpha\beta) - \bar{r}(\alpha\beta) = \bar{d}(\alpha) - \bar{r}(\alpha) + \bar{d}(\beta) - \bar{r}(\beta).$$

(2) Suppose that the case when $\text{dom } \beta \subseteq \text{ran } \alpha$ holds. We put $k = \bar{d}(\beta) - \bar{r}(\alpha)$. Then the definition of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ implies that $k \geq 0$, $\bar{d}(\alpha\beta) = \bar{d}(\alpha) + k$, $\bar{r}(\alpha\beta) = \bar{r}(\beta)$, and hence in this case we have that

$$\bar{d}(\alpha\beta) - \bar{r}(\alpha\beta) = \bar{d}(\alpha) - \bar{r}(\alpha) + \bar{d}(\beta) - \bar{r}(\beta).$$

(3) Now we consider the case $(\lambda \setminus \text{ran } \alpha) \cap (\lambda \setminus \text{dom } \beta) \neq \emptyset$, $\text{ran } \alpha \not\subseteq \text{dom } \beta$ and $\text{dom } \beta \not\subseteq \text{ran } \alpha$. Then the definition of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ implies that there exist positive integers i , j and k such that $i = |(\lambda \setminus \text{ran } \alpha) \setminus (\lambda \setminus \text{dom } \beta)|$, $j = |(\lambda \setminus \text{ran } \alpha) \cap (\lambda \setminus \text{dom } \beta)|$ and $k = |(\lambda \setminus \text{dom } \beta) \setminus (\lambda \setminus \text{ran } \alpha)|$. Then we have that $\bar{r}(\alpha) = i + j$, $\bar{d}(\beta) = j + k$, $\bar{d}(\alpha\beta) = \bar{d}(\alpha) + k$ and $\bar{r}(\alpha\beta) = \bar{r}(\beta) + i$. Therefore, in this case we get that

$$\bar{d}(\alpha\beta) - \bar{r}(\alpha\beta) = \bar{d}(\alpha) - \bar{r}(\alpha) + \bar{d}(\beta) - \bar{r}(\beta).$$

(4) In the case when $(\lambda \setminus \text{ran } \alpha) \cap (\lambda \setminus \text{dom } \beta) = \emptyset$ we have that the definition of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ implies that $\bar{d}(\alpha\beta) = \bar{d}(\alpha) + \bar{d}(\beta)$, $\bar{r}(\alpha\beta) = \bar{r}(\alpha) + \bar{r}(\beta)$, and hence we get that

$$\bar{d}(\alpha\beta) - \bar{r}(\alpha\beta) = \bar{d}(\alpha) - \bar{r}(\alpha) + \bar{d}(\beta) - \bar{r}(\beta).$$

This completes the proof of the proposition. \square

On the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ we define a relation $\sim_{\mathfrak{d}}$ in the following way:

$$\alpha \sim_{\mathfrak{d}} \beta \quad \text{if and only if} \quad \bar{d}(\alpha) - \bar{r}(\alpha) = \bar{d}(\beta) - \bar{r}(\beta),$$

for $\alpha, \beta \in \mathcal{J}_\lambda^{\text{cf}}$.

Proposition 3.3. *Let λ be an infinite cardinal. Then $\sim_{\mathfrak{d}}$ is a congruence on the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$ and moreover the quotient semigroup $\mathcal{J}_{\lambda}^{\text{cf}} / \sim_{\mathfrak{d}}$ is isomorphic to the additive group of integers $\mathbb{Z}(+)$.*

Proof. Simple verifications and Proposition 3.2 imply that $\sim_{\mathfrak{d}}$ is a congruence on the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$. We define a homomorphism $h: \mathcal{J}_{\lambda}^{\text{cf}} \rightarrow \mathbb{Z}(+)$ by the formula $(\alpha)h = \bar{d}(\alpha) - \bar{r}(\alpha)$. Then the definitions of the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$ and the congruence $\sim_{\mathfrak{d}}$ on $\mathcal{J}_{\lambda}^{\text{cf}}$, and Proposition 3.2 imply that thus defined map h is a surjective homomorphism and moreover $(\alpha)h = (\beta)h$ if and only if $\alpha \sim_{\mathfrak{d}} \beta$ in $\mathcal{J}_{\lambda}^{\text{cf}}$. This completes the proof of the proposition. \square

Proposition 3.4. *Let λ be an infinite cardinal. Then for every element β of the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$ such that $\bar{d}(\beta) = \bar{r}(\beta)$ there exists an element α of the group of units of $\mathcal{J}_{\lambda}^{\text{cf}}$ such that $\alpha\sigma\beta$.*

Proof. Fix an arbitrary element β of the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$. Without loss of generality we can assume that $\bar{d}(\beta) = \bar{r}(\beta) = k > 0$. Let $\{x_1, \dots, x_k\} = \lambda \setminus \text{dom } \beta$ and $\{y_1, \dots, y_k\} = \lambda \setminus \text{ran } \beta$. We define a map $\alpha: \lambda \rightarrow \lambda$ in the following way:

$$(x)\alpha = \begin{cases} (x)\beta, & \text{if } x \in \text{dom } \beta; \\ y_i, & \text{if } x = x_i, i = 1, \dots, k. \end{cases}$$

Then α is an element of the group of units of the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$ and it is obviously that $\alpha\varepsilon = \beta\varepsilon$, where ε is the identity map of the set $\text{ran } \beta$. \square

For every $\alpha \in \mathcal{S}_{\lambda}$ we denote $\text{supp}(\alpha) = \{x \in \lambda \mid (x)\alpha \neq x\}$. We define

$$\mathcal{S}_{\lambda}^{\infty} = \{\alpha \in \mathcal{S}_{\lambda} \mid \text{supp}(\alpha) \text{ is finite}\}.$$

We observe that the Schreier–Ulam theorem (see [14, Theorem 11.3.4]) implies that $\mathcal{S}_{\lambda}^{\infty}$ is a normal subgroup of \mathcal{S}_{λ} and hence $\mathcal{S}_{\lambda}/\mathcal{S}_{\lambda}^{\infty}$ is a group.

Later on, when \mathfrak{C} is a congruence on a semigroup S we shall denote the *natural homomorphism* generated by the congruence \mathfrak{C} on S by $\pi_{\mathfrak{C}}: S \rightarrow S/\mathfrak{C}$.

The definition of the least group congruence σ on the semigroup $\mathcal{J}_{\lambda}^{\text{cf}}$ implies the following proposition.

Proposition 3.5. *Let λ be an infinite cardinal. Then the homomorphic image $(H(\mathbb{I}))\pi_{\sigma}$ of the group of units $H(\mathbb{I})$ of $\mathcal{J}_{\lambda}^{\text{cf}}$ under the natural homomorphism $\pi_{\sigma}: \mathcal{J}_{\lambda}^{\text{cf}} \rightarrow \mathcal{J}_{\lambda}^{\text{cf}}/\sigma$ is isomorphic to the quotient group $\mathcal{S}_{\lambda}/\mathcal{S}_{\lambda}^{\infty}$.*

Theorem 3.6. *Let λ be an infinite cardinal. Then the following conditions hold:*

- (i) $(H(\mathbb{I}))\pi_{\sigma} = \mathcal{S}_{\lambda}/\mathcal{S}_{\lambda}^{\infty}$ is a normal subgroup of the group $\mathcal{J}_{\lambda}^{\text{cf}}/\sigma$;
- (ii) The group $\mathcal{J}_{\lambda}^{\text{cf}}/\sigma$ contains the infinite cyclic subgroup G (i.e., the additive group of integers $\mathbb{Z}(+)$) such that $G \cap \mathcal{S}_{\lambda}/\mathcal{S}_{\lambda}^{\infty} = \{e\}$, where e is the unit of the group $\mathcal{J}_{\lambda}^{\text{cf}}/\sigma$;
- (iii) $\mathcal{S}_{\lambda}/\mathcal{S}_{\lambda}^{\infty} \cdot G = \mathcal{J}_{\lambda}^{\text{cf}}/\sigma$.

and hence the group $\mathcal{J}_{\lambda}^{\text{cf}}/\sigma$ is isomorphic to the semidirect product $\mathcal{S}_{\lambda}/\mathcal{S}_{\lambda}^{\infty} \rtimes \mathbb{Z}(+)$.

Proof. (i) Since $\sigma \subseteq \sim_{\mathfrak{d}}$ we conclude that Theorem 1.6 of [5] implies that there exists a unique homomorphism $g: \mathcal{J}_{\lambda}^{\text{cf}}/\sigma \rightarrow G$ such that the following diagram

$$\begin{array}{ccc} \mathcal{J}_{\lambda}^{\text{cf}} & \xrightarrow{\pi_{\sigma}} & \mathcal{J}_{\lambda}^{\text{cf}}/\sigma \\ & \searrow \pi_{\sim_{\mathfrak{d}}} & \downarrow g \\ & & G \end{array}$$

commutes. Then by Proposition 3.5 we have that the homomorphic image $(H(\mathbb{I}))\pi_\sigma$ of the group of units $H(\mathbb{I})$ of $\mathcal{J}_\lambda^{\text{cf}}$ under the natural homomorphism $\pi_\sigma: \mathcal{J}_\lambda^{\text{cf}} \rightarrow \mathcal{J}_\lambda^{\text{cf}}/\sigma$ is isomorphic to the quotient group $\mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty$. Now Propositions 3.4 and 3.5 imply that the subgroup $(H(\mathbb{I}))\pi_\sigma = \mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty$ of the group $\mathcal{J}_\lambda^{\text{cf}}/\sigma$ is the kernel of the homomorphism $g: \mathcal{J}_\lambda^{\text{cf}}/\sigma \rightarrow G$, and hence $(H(\mathbb{I}))\pi_\sigma = \mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty$ is a normal subgroup of $\mathcal{J}_\lambda^{\text{cf}}/\sigma$.

(ii) Fix an arbitrary $\alpha \in \mathcal{J}_\lambda^{\text{cf}}$ such that $|\lambda \setminus \text{dom } \alpha| = 1$ and $\text{ran } \alpha = \lambda$. Then the definition of the congruence \sim_∂ on $\mathcal{J}_\lambda^{\text{cf}}$ implies that the element α^n is not \sim_∂ -equivalent to any element of the group of units $H(\mathbb{I})$ for every non-zero integer n , and hence by Proposition 3.5 we get that $((\alpha)\pi_\sigma)^n \notin \mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty$. This implies that $\{((\alpha)\pi_\sigma)^n \mid n \in \mathbb{Z}\} \cap \mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty = \{e\}$, where e is the unit of the group $\mathcal{J}_\lambda^{\text{cf}}/\sigma$. Also, it is obvious that $(\alpha^n)\pi_\sigma = n \in G$ and $\{(\alpha^n)\pi_\sigma \mid n \in \mathbb{Z}\}$ is a cyclic subgroup of $\mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty$.

(iii) Fix an arbitrary element x in $\mathcal{J}_\lambda^{\text{cf}}/\sigma$. Let ξ be an arbitrary element of the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ be such that $(\xi)\pi_\sigma = x$. If $\bar{d}(\xi) = \bar{r}(\xi)$ then by Proposition 3.4 we have that $\xi\sigma\beta$ for some element β from the group of units of $\mathcal{J}_\lambda^{\text{cf}}$, and hence we get that $x = (\beta)\pi_\sigma \cdot e \in \mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty \cdot G$, where e is the unit of the group $\mathcal{J}_\lambda^{\text{cf}}/\sigma$. Suppose that $\bar{d}(\xi) - \bar{r}(\xi) = n \neq 0$. Then by Proposition 3.2 we have that $\bar{d}(\xi \cdot (\alpha^{-1})^n) - \bar{r}(\xi \cdot (\alpha^{-1})^n) = 0$. Now, Proposition 3.4 implies that the element $\xi \cdot (\alpha^{-1})^n$ is σ -equivalent to some element β of the group of units $H(\mathbb{I})$ of $\mathcal{J}_\lambda^{\text{cf}}$. Then we have that $(\xi \cdot (\alpha^{-1})^n)\pi_\sigma = (\beta)\pi_\sigma$ and since $\mathcal{J}_\lambda^{\text{cf}}/\sigma$ is a group we get that $x = (\xi)\pi_\sigma = (\beta)\pi_\sigma \cdot (\alpha^n)\pi_\sigma \in \mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty \cdot G$. This implies that $\mathcal{S}_\lambda/\mathcal{S}_\lambda^\infty \cdot G = \mathcal{J}_\lambda^{\text{cf}}/\sigma$.

The last statement of the theorem follows from statements (i)–(iii) and Exercise 2.5.3 from [6]. \square

Remark 3.7. *Proposition 3.3 implies that for every infinite cardinal λ the group $\mathcal{J}_\lambda^{\text{cf}}/\sigma$ has infinitely many normal subgroups and hence the semigroup $\mathcal{J}_\lambda^{\text{cf}}$ has infinitely many group congruences.*

Acknowledgements. This research was supported by the Slovenian Research Agency grants P1-0292, J1-5435, and J1-6721. The authors are grateful to the referee for several useful comments and suggestions.

REFERENCES

- [1] ANDERSEN, O.: *Ein Bericht über die Struktur abstrakter Halbgruppen*, PhD Thesis, Hamburg, 1952.
- [2] ANDERSON, L. W.—HUNTER, R. P.—KOCH, R. J.: *Some results on stability in semigroups*, Trans. Amer. Math. Soc. **117** (1965), 521–529.
- [3] BANAKH, T.—DIMITROVA, S.—GUTIK, O.: *The Rees-Suschkiewitsch theorem for simple topological semigroups*, Mat. Stud. **31** (2009), 211–218.
- [4] BANAKH, T.—DIMITROVA, S.—GUTIK, O.: *Embedding the bicyclic semigroup into countably compact topological semigroups*, Topology Appl. **157** (2010), 2803–2814.
- [5] CLIFFORD, A. H.—PRESTON, G. B.: *The Algebraic Theory of Semigroups*, Vol. I/II, American Mathematical Society, Providence, Rhode Island, 1961/1967.
- [6] DIXON J. D.—MORTIMER B.: *Permutation Groups*, Springer, Berlin, 1996.
- [7] EBERHART, C.—SELDEN, J.: *On the closure of the bicyclic semigroup*, Trans. Amer. Math. Soc. **144** (1969), 115–126.
- [8] GUTIK, O.—REPOVŠ, D.: *On countably compact 0-simple topological inverse semigroups*, Semigroup Forum **75** (2007), 464–469.

- [9] GUTIK, O.—REPOVŠ, D.: *Topological monoids of monotone injective partial selfmaps of N with cofinite domain and image*, Stud. Sci. Math. Hung. **48** (2011), 342–353.
- [10] GUTIK, O.—REPOVŠ, D.: *On monoids of injective partial selfmaps of integers with cofinite domains and images*, Georgian Math. J. **19** (2012), 511–532.
- [11] HILDEBRANT, J. A.—KOCH, R. J.: *Swelling actions of Γ -compact semigroups*, Semigroup Forum **33** (1986), 65–85.
- [12] HOWIE, J. M.: *Fundamentals of Semigroup Theory*, London Math. Soc. Monogr. (N.S.) No. 12, Clarendon Press, Oxford, 1995.
- [13] PETRICH, M.: *Inverse Semigroups*, Pure Appl. Math., John Wiley & Sons, New York, 1984.
- [14] SCOTT, W. R.: *Group Theory*, Dover Publications, Inc., New York, 1987.
- [15] SHELAH, S.—STEPRĀNS, J.: *Non-trivial homeomorphisms of $\beta N \setminus N$ without the Continuum Hypothesis*, Fund. Math. **132** (1989), 135–141.
- [16] SHELAH, S.—STEPRĀNS, J.: *Somewhere trivial autohomeomorphisms*, J. London Math. Soc. (2), **49** (1994), 569–580.
- [17] SHELAH, S.—STEPRĀNS, J.: *Martin's axiom is consistent with the existence of nowhere trivial automorphisms*, Proc. Amer. Math. Soc. **130** (2002), 2097–2106.
- [18] VELIČKOVIĆ, B.: *Definable automorphisms of $\mathcal{P}(\omega)/\text{fin}$* , Proc. Amer. Math. Soc. **96** (1986), 130–135.
- [19] VELIČKOVIĆ, B.: *Applications of the Open Coloring Axiom*, In Set Theory of the Continuum, H. Judah, W. Just et H. Woodin, eds., Pap. Math. Sci. Res. Inst. Workshop, Berkeley, 1989, MSRI Publications. Springer-Verlag. Vol. **26**, Berlin, (1992), pp. 137–154.
- [20] VELIČKOVIĆ, B.: *OCA and automorphisms of $\mathcal{P}(\omega)/\text{fin}$* , Topology Appl. **49** (1993), 1–13.
- [21] VAGNER, V. V.: *Generalized groups*, Dokl. Akad. Nauk SSSR **84** (1952), 1119–1122 (in Russian).

DEPARTMENT OF MECHANICS AND MATHEMATICS
 IVAN FRANKO NATIONAL UNIVERSITY OF LVIV
 UNIVERSYTETSKA 1
 LVIV, 79000
 UKRAINE
E-mail address: `o.gutik@franko.lviv.ua`

FACULTY OF EDUCATION, AND
 FACULTY OF MATHEMATICS AND PHYSICS
 UNIVERSITY OF LJUBLJANA
 KARDELJEVA PLOŠČAD 16
 LJUBLJANA, 1000
 SLOVENIA
E-mail address: `dusan.repovs@guest.arnes.si`