

EXISTENCE AND MULTIPLICITY OF SOLUTIONS FOR DOUBLE-PHASE ROBIN PROBLEMS

N.S. PAPAGEORGIOU, V.D. RĂDULESCU, AND D.D. REPOVŠ

ABSTRACT. We consider a double phase Robin problem with a Carathéodory nonlinearity. When the reaction is superlinear but without satisfying the Ambrosetti-Rabinowitz condition, we prove an existence theorem. When the reaction is resonant, we prove a multiplicity theorem. Our approach is Morse theoretic, using the notion of homological local linking.

1. INTRODUCTION

Let $\Omega \subseteq \mathbb{R}^N$ be a bounded domain with Lipschitz boundary $\partial\Omega$. In this paper we study the following two phase Robin problem

$$(1) \quad \left\{ \begin{array}{ll} -\operatorname{div}(a_0(z)|Du|^{p-2}Du) - \Delta_q u + \xi(z)|u|^{p-2}u = f(z, u) & \text{in } \Omega \\ \frac{\partial u}{\partial n_\theta} + \beta(z)|u|^{p-2}u = 0 & \text{on } \partial\Omega, \end{array} \right\}$$

where $1 < q < p \leq N$.

In this problem, the weight $a_0 : \bar{\Omega} \rightarrow \mathbb{R}$ is Lipschitz continuous and $a_0(z) > 0$ for all $z \in \Omega$. The potential function $\xi \in L^\infty(\Omega)$ satisfies $\xi(z) \geq 0$ for a.a. $z \in \Omega$, while the reaction term $f(z, x)$ is Carathéodory (that is, for all $x \in \mathbb{R}$ the mapping $z \mapsto f(z, x)$ is measurable and for a.a. $z \in \Omega$ the function $x \mapsto f(z, x)$ is continuous). Let $F(z, \cdot)$ be the primitive of $f(z, \cdot)$, that is, $F(z, x) = \int_0^x f(z, s)ds$. We assume that for a.a. $z \in \Omega$, $F(z, \cdot)$ is q -linear near the origin. On the other hand, near $\pm\infty$, we consider two distinct cases for $f(z, \cdot)$:

- (i) for a.a. $z \in \Omega$, $f(z, \cdot)$ is $(p-1)$ -superlinear but without satisfying the Ambrosetti-Rabinowitz condition (the AR-condition for short), which is common in the literature when dealing with super-linear problems;
- (ii) for a.a. $z \in \Omega$, $f(z, \cdot)$ is $(p-1)$ -linear and possibly resonant with respect to the principal eigenvalue of the weighted p -Laplacian

$$u \mapsto -\operatorname{div}(a_0(z)|Du|^{p-2}Du)$$

with Robin boundary condition.

In the boundary condition, $\frac{\partial u}{\partial n_\theta}$ denotes the conormal derivative of u corresponding to the modular function $\theta(z, x) = a_0(z)x^p + x^q$ for all $z \in \Omega$, all $x \geq 0$. We interpret this derivative via the nonlinear Green identity (see Papageorgiou, Rădulescu and Repovš [18, p. 34]) and

$$\frac{\partial u}{\partial n_\theta} = [a_0(z)|Du|^{p-2} + |Du|^{q-2}] \frac{\partial u}{\partial n} \text{ for all } u \in C^1(\bar{\Omega}),$$

with $n(\cdot)$ being the outward unit normal on $\partial\Omega$. The boundary coefficient β satisfies $\beta \in C^{0,\alpha}(\partial\Omega)$ with $0 < \alpha < 1$ and $\beta(z) \geq 0$ for all $z \in \partial\Omega$.

The differential operator in problem (1) is a weighted (p, q) -Laplace operator and it corresponds to the energy functional

$$u \mapsto \int_{\Omega} [a_0(z)|Du|^p + |Du|^q] dz.$$

Key words and phrases. Unbalanced growth, Musielak-Orlicz space, homological local linking, superlinear reaction, resonant reaction.

2010 Mathematics Subject Classification. Primary: 35J20. Secondary: 35J25, 35J60.

Since we do not assume that the weight function $a_0(z)$ is bounded away from zero, the continuous integrand $\theta_0 : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}_+$ of this integral functional exhibits unbalanced growth, namely

$$|y|^q \leq \theta_0(z, y) \leq c_0(1 + |y|^p) \text{ for a.a. } z \in \Omega, \text{ all } y \in \mathbb{R}^N \text{ and some } c_0 > 0.$$

Such integral functionals were first investigated by Marcellini [14] and Zhikov [22], in connection with problems in nonlinear elasticity theory. Recently, Baroni, Colombo and Mingione [3] and Colombo and Mingione [6, 7] revived the interest in them and produced important local regularity results for the minimizers of such functionals. A global regularity theory for such problems remains elusive.

In this paper, using tools from Morse theory (in particular, critical groups), we prove an existence theorem (for the superlinear case) and a multiplicity theorem (for the linear resonant case). Existence and multiplicity results for two phase problems were proved recently by Cencelj, Rădulescu and Repovš [4] (problems with variable growth), Colasuonno and Squassina [5] (eigenvalue problems), Liu and Dai [13] (existence of solutions for problems with superlinear reaction), Papageorgiou, Rădulescu and Repovš [19] (multiple solutions for superlinear problems), and Papageorgiou, Vetro and Vetro [20] (parametric Dirichlet problems). The approach in all the aforementioned works is different and the hypotheses on the reaction are more restrictive.

Finally, we mention that (p, q) -equations arise in many mathematical models of physical processes. We refer to the very recent works of Bahrouni, Rădulescu and Repovš [1, 2] and the references therein.

2. MATHEMATICAL BACKGROUND

The study of two-phase problems requires the use of Musielak-Orlicz spaces. So, let $\theta : \Omega \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be the modular function defined by

$$\theta(z, x) = a_0(z)x^p + x^q \text{ for all } z \in \Omega, x \geq 0.$$

This is a generalized N-function (see Musielak [16]) and it satisfies

$$\theta(z, 2x) \leq 2^p \theta(z, x) \text{ for all } z \in \Omega, x \geq 0,$$

that is, $\theta(z, \cdot)$ satisfies the (Δ_2) -property (see Musielak [16, p. 52]). Using the modular function $\theta(z, x)$, we can define the Musielak-Orlicz space $L^\theta(\Omega)$ as follows:

$$L^\theta(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R}; u \text{ is measurable and } \int_\Omega \theta(z, |u|) dz < \infty \right\}.$$

This space is equipped with the so-called ‘‘Luxemburg norm’’ defined by

$$\|u\|_\theta = \inf \left\{ \lambda > 0 : \int_\Omega \theta(z, \frac{|u|}{\lambda}) dz \leq 1 \right\}.$$

Using $L^\theta(\Omega)$, we can define the following Sobolev-type space $W^{1,\theta}(\Omega)$, by setting

$$W^{1,\theta}(\Omega) = \{u \in L^\theta(\Omega) : |Du| \in L^\theta(\Omega)\}.$$

We equip $W^{1,\theta}(\Omega)$ with the norm $\|\cdot\|$ defined by

$$\|u\| = \|u\|_\theta + \|Du\|_\theta,$$

where $\|Du\|_\theta = \||Du|\|_\theta$. The spaces $L^\theta(\Omega)$ and $W^{1,\theta}(\Omega)$ are separable and uniformly convex (hence reflexive) Banach spaces.

Let $\hat{\theta}(z, x)$ be another modular function. We say that ‘‘ $\hat{\theta}$ is weaker than θ ’’ and write $\hat{\theta} \prec \theta$, if there exist $c_1, c_2 > 0$ and a function $\eta \in L^1(\Omega)$ such that

$$\hat{\theta}(z, x) \leq c_1 \theta(z, c_2 x) + \eta(z) \text{ for a.a. } z \in \Omega \text{ and all } x \geq 0.$$

Then we have

$$L^\theta(\Omega) \hookrightarrow L^{\hat{\theta}}(\Omega) \text{ and } W^{1,\theta}(\Omega) \hookrightarrow W^{1,\hat{\theta}}(\Omega) \text{ continuously.}$$

Combining this fact with the classical Sobolev embedding theorem, we obtain the following embeddings; see Propositions 2.15 and 2.18 of Colasuonno and Squassina [5].

Proposition 2.1. *We assume that $1 < q < p < \infty$. Then the following properties hold.*

(a) *If $q \neq N$, then $W^{1,\theta}(\Omega) \hookrightarrow L^r(\Omega)$ continuously for all $1 \leq r \leq q^*$, where*

$$q^* = \begin{cases} \frac{Nq}{N-q} & \text{if } q < N \\ +\infty & \text{if } q \geq N. \end{cases}$$

(b) *If $q = N$, then $W^{1,\theta}(\Omega) \hookrightarrow L^r(\Omega)$ continuously for all $1 \leq r < \infty$.*

(c) *If $q \leq N$, then $W^{1,\theta}(\Omega) \hookrightarrow L^r(\Omega)$ compactly for all $1 \leq r < q^*$.*

(d) *If $q > N$, then $W^{1,\theta}(\Omega) \hookrightarrow L^\infty(\Omega)$ compactly.*

(e) *$W^{1,\theta}(\Omega) \hookrightarrow W^{1,q}(\Omega)$ continuously.*

We have

$$L^p(\Omega) \hookrightarrow L^\theta(\Omega) \hookrightarrow L_{a_0}^p(\Omega) \cap L^q(\Omega)$$

with both embeddings being continuous.

We consider the modular function

$$\rho_\theta(u) = \int_{\Omega} \theta(z, |Du|) dz = \int_{\Omega} [a_0(z)|Du|^p + |Du|^q] dz \text{ for all } u \in W^{1,\theta}(\Omega).$$

There is a close relationship between the norm $\|\cdot\|$ of $W^{1,\theta}(\Omega)$ and the modular functional $\rho_\theta(\cdot)$; see Proposition 2.1 of Liu and Dai [13].

Proposition 2.2. (a) *If $u \neq 0$, then $\|Du\|_\theta = \lambda$ if and only if $\rho_\theta(\frac{u}{\lambda}) \leq 1$.*

(b) *$\|Du\|_\theta < 1$ (resp. $= 1, > 1$) if and only if $\rho_\theta(u) < 1$ (resp. $= 1, > 1$).*

(c) *If $\|Du\|_\theta < 1$, then $\|Du\|_\theta^p \leq \rho_\theta(u) \leq \|Du\|_\theta^q$.*

(d) *If $\|Du\|_\theta > 1$, then $\|Du\|_\theta^q \leq \rho_\theta(u) \leq \|Du\|_\theta^p$.*

(e) *$\|Du\|_\theta \rightarrow 0$ if and only if $\rho_\theta(u) \rightarrow 0$.*

(f) *$\|Du\|_\theta \rightarrow +\infty$ if and only if $\rho_\theta(u) \rightarrow +\infty$.*

On $\partial\Omega$ we consider the $(N-1)$ -dimensional Hausdorff (surface) measure $\sigma(\cdot)$. Using this measure, we can define in the usual way the ‘‘boundary’’ Lebesgue spaces $L^s(\partial\Omega)$ for $1 \leq s \leq \infty$. It is well-known that there exists a unique continuous linear map $\gamma_0 : W^{1,q}(\Omega) \rightarrow L^q(\partial\Omega)$, known as the ‘‘trace map’’, such that

$$\gamma_0(u) = u|_{\partial\Omega} \text{ for all } u \in W^{1,q}(\Omega) \cap C(\bar{\Omega}).$$

We have

$$\text{im } \gamma_0 = W^{\frac{1}{q},q}(\Omega) \left(\frac{1}{q} + \frac{1}{q'} = 1 \right) \text{ and } \ker \gamma_0 = W_0^{1,q}(\Omega).$$

Moreover, the trace map $\gamma_0(\cdot)$ is compact into $L^s(\partial\Omega)$ for all $1 \leq s < (N-1)q/(N-q)$ if $q < N$, and for all $1 \leq s < \infty$ if $q \geq N$. In what follows, for the sake of notational simplicity, we drop the use of the trace map $\gamma_0(\cdot)$. All restrictions of the Sobolev functions on the boundary $\partial\Omega$ are understood in the sense of traces.

Let $\langle \cdot, \cdot \rangle$ denote the duality brackets for the pair $(W^{1,\theta}(\Omega), W^{1,\theta}(\Omega)^*)$ and $\langle \cdot, \cdot \rangle_{1,q}$ denote the duality brackets for the pair $(W^{1,q}(\Omega), W^{1,q}(\Omega)^*)$. We introduce the maps $A_p^{a_0} : W^{1,\theta}(\Omega) \rightarrow W^{1,\theta}(\Omega)^*$ and $A_q : W^{1,q}(\Omega) \rightarrow W^{1,q}(\Omega)^*$ defined by

$$\langle A_p^{a_0}(u), h \rangle = \int_{\Omega} a_0(z) |Du|^{p-2} (Du, Dh)_{\mathbb{R}^N} dz \text{ for all } u, h \in W^{1,\theta}(\Omega),$$

$$\langle A_q(u), h \rangle_{1,q} = \int_{\Omega} |Du|^{q-2} (Du, Dh)_{\mathbb{R}^N} dz \text{ for all } u, h \in W^{1,q}(\Omega).$$

We have

$$\langle A_q(u), h \rangle_{1,q} = \langle A_q(u), h \rangle \text{ for all } u, h \in W^{1,\theta}(\Omega).$$

We introduce the following hypotheses on the weight $a_0(\cdot)$ and on the coefficients $\xi(\cdot)$ and $\beta(\cdot)$.

H_0 : $a_0 : \bar{\Omega} \rightarrow \mathbb{R}$ is Lipschitz continuous, $a_0(z) > 0$ for all $z \in \Omega$, $\xi \in L^\infty(\Omega)$, $\xi(z) \geq 0$ for a.a. $z \in \Omega$, $\beta \in C^{0,\alpha}(\partial\Omega)$ with $0 < \alpha < 1$, $\xi \not\equiv 0$ or $\beta \not\equiv 0$ and $q > Np/(N+p-1)$.

Remark 2.1. *The latter condition on the exponent q implies that $W^{1,\theta}(\Omega) \hookrightarrow L^p(\partial\Omega)$ compactly and $q < p^*$.*

We introduce the C^1 -functional $\gamma_p : W^{1,\theta}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\gamma_p(u) = \int_{\Omega} a_0(z)|Du|^p dz + \int_{\Omega} \xi(z)|u|^p dz + \int_{\partial\Omega} \beta(z)|u|^p d\sigma \text{ for all } u \in W^{1,\theta}(\Omega).$$

Then hypotheses H_0 , Lemma 4.11 of Mugnai and Papageorgiou [15], and Proposition 2.4 of Gasinski and Papageorgiou [10], imply that

$$(2) \quad c_1 \|u\|^p \leq \gamma_p(u) \text{ for some } c_1 > 0, \text{ all } u \in W^{1,\theta}(\Omega).$$

We denote by $\hat{\lambda}_1(p)$ the first (principal) eigenvalue of the following nonlinear eigenvalue problem

$$(3) \quad \left\{ \begin{array}{ll} -\operatorname{div}(a_0(z)|Du|^{p-2}Du) + \xi(z)|u|^{p-2}u = \hat{\lambda}|u|^{p-2}u & \text{in } \Omega \\ \frac{\partial u}{\partial n_p} + \beta(z)|u|^{p-2}u = 0 & \text{on } \partial\Omega. \end{array} \right.$$

Here, $\frac{\partial u}{\partial n_p} = |Du|^{p-2} \frac{\partial u}{\partial n}$. The eigenvalue $\hat{\lambda}_1(p)$ has the following variational characterization

$$(4) \quad \hat{\lambda}_1(p) = \inf \left\{ \frac{\gamma_p(u)}{\|u\|_p^p} : u \in W^{1,p}(\Omega) \setminus \{0\} \right\} \text{ (see [17]).}$$

Then by (2), we see that $\hat{\lambda}_1(p) > 0$. This eigenvalue is simple (that is, if \hat{u}, \hat{v} are corresponding eigenfunctions, then $\hat{u} = \eta\hat{v}$ with $\eta \in \mathbb{R} \setminus \{0\}$) and isolated (that is, if $\hat{\sigma}(p)$ denotes the spectrum of (3), then we can find $\varepsilon > 0$ such that $(\hat{\lambda}_1(p), \hat{\lambda}_1(p) + \varepsilon) \cap \hat{\sigma}(p) = \emptyset$). The infimum in (4) is realized on the corresponding one-dimensional eigenspace, the elements of which have fixed sign. We denote by $\hat{u}_1(p)$ the corresponding positive, L^p -normalized (that is, $\|\hat{u}_1(p)\|_p = 1$) eigenfunction. We know that $\hat{u}_1(p) \in L^\infty(\Omega)$ (see Colasuonno and Squassina [5, Section 3.2]) and $\hat{u}_1(p)(z) > 0$ for a.a. $z \in \Omega$ (see Papageorgiou, Vetro and Vetro [19, Proposition 4]).

We will also use the spectrum of the following nonlinear eigenvalue problem

$$-\Delta_q u = \hat{\lambda}|u|^{q-2}u \text{ in } \Omega, \quad \frac{\partial u}{\partial n} = 0 \text{ on } \partial\Omega.$$

It is well known that this problem has a sequence of variational eigenvalues $\{\hat{\lambda}_k(q)\}_{k \geq 1}$ such that $\hat{\lambda}_k(q) \rightarrow +\infty$ as $k \rightarrow \infty$. We have $\hat{\lambda}_1(q) = 0 < \hat{\lambda}_2(q)$ (see Gasinski and Papageorgiou [9, Section 6.2]).

Let X be a Banach space and $\phi \in C^1(X, \mathbb{R})$. We denote by K_ϕ the critical set of ϕ , that is,

$$K_\phi = \{u \in X : \phi'(u) = 0\}.$$

Also, if $\eta \in \mathbb{R}$, then we set

$$\phi^\eta = \{u \in X : \phi(u) \leq \eta\}.$$

Consider a topological pair (A, B) such that $B \subseteq A \subseteq X$. Then for every $k \in \mathbb{N}_0$, we denote by $H_k(A, B)$ the k th-singular homology group for the pair (A, B) with coefficients in a field \mathbb{F} of characteristic zero (for example, $\mathbb{F} = \mathbb{R}$). Then each $H_k(A, B)$ is an \mathbb{F} -vector space and we denote by $\dim H_k(A, B)$ its dimension. We also recall that the homeomorphisms induced by maps of pairs and the boundary homomorphism ∂ , are all \mathbb{F} -linear.

Suppose that $u \in K_\phi$ is isolated. Then for every $k \in \mathbb{N}_0$, we define the “ k -critical group” of ϕ at u by

$$C_k(\phi, u) = H_k(\phi^c \cap U, \phi^c \cap U \setminus \{u\}),$$

where U is an isolating neighborhood of u , that is, $K_\phi \cap U \cap \phi^c = \{u\}$. The excision property of singular homology implies that this definition is independent of the choice of the isolating neighborhood U .

We say that ϕ satisfies the “C-condition” if it has the following property:

“Every sequence $\{u_n\}_{n \geq 1} \subseteq X$ such that $\{\phi(u_n)\}_{n \geq 1} \subseteq \mathbb{R}$ is bounded and $(1 + \|u_n\|)\phi'(u_n) \rightarrow 0$ in X^* as $n \rightarrow \infty$, has a strongly convergent subsequence”.

Suppose that $\phi \in C^1(X, \mathbb{R})$ satisfies the C-condition and that $\inf \phi(K_\phi) > -\infty$. Let $c < \inf \phi(K_\phi)$. Then the critical groups of ϕ at infinity are defined by

$$C_k(\phi, \infty) = H_k(X, \phi^c) \text{ for all } k \in \mathbb{N}_0.$$

On account of the second deformation theorem (see Papageorgiou, Rădulescu and Repovš [18, p. 386], Theorem 5.3.12) this definition is independent of the choice of the level $c < \inf \phi(K_\phi)$.

Our approach is based on the notion of local (m, n) -linking ($m, n \in \mathbb{N}$), see Papageorgiou, Rădulescu and Repovš [18, Definition 6.6.13, p. 534].

Definition 2.3. *Let X be a Banach space, $\phi \in C^1(X, \mathbb{R})$, and 0 an isolated critical point of ϕ with $\phi(0) = 0$. Let $m, n \in \mathbb{N}$. We say that ϕ has a “local (m, n) -linking” near the origin if there exist a neighborhood U of 0 and nonempty sets $E_0, E \subseteq U$, and $D \subseteq X$ such that $0 \notin E_0 \subseteq E$, $E_0 \cap D = \emptyset$ and*

(a) 0 is the only critical point of ϕ in $\phi^0 \cap U$;

(b) $\dim \operatorname{im} i_* - \dim \operatorname{im} j_* \geq n$, where

$$i_* : H_{m-1}(E_0) \rightarrow H_{m-1}(X \setminus D) \text{ and } j_* : H_{m-1}(E_0) \rightarrow H_{m-1}(E)$$

are the homomorphisms induced by the inclusion maps $i : E_0 \rightarrow X \setminus D$ and $j : E_0 \rightarrow E$;

(c) $\phi|_E \leq 0 < \phi|_{U \cap D \setminus \{0\}}$.

Remark 2.2. *The notion of “local (m, n) -linking” was introduced by Perera [21] as a generalization of the concept of local linking due to Liu [12]. Here we introduce a slightly more general version of this notion.*

3. SUPERLINEAR CASE

In this section we treat the superlinear case, that is, we assume that the reaction $f(z, \cdot)$ exhibits $(p-1)$ -superlinear growth near $\pm\infty$.

The hypotheses on $f(z, x)$ are the following.

H_1 : $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function such that $f(z, 0) = 0$ for a.a. $z \in \Omega$ and

(i) $|f(z, x)| \leq \hat{a}(z)(1 + |x|^{r-1})$ for a.a. $z \in \Omega$ and all $x \in \Omega$, with $\hat{a} \in L^\infty(\Omega)$, $p < r < q^*$;

(ii) if $F(z, x) = \int_0^x f(z, s) ds$, then $\lim_{x \rightarrow \pm\infty} \frac{F(z, x)}{|x|^p} = +\infty$ uniformly for a.a. $z \in \Omega$;

(iii) if $\eta(z, x) = f(z, x)x - pF(z, x)$, then there exists $e \in L^1(\Omega)$ such that

$$\eta(z, x) \leq \eta(z, y) + e(z) \text{ for a.a. } z \in \Omega \text{ and all } 0 \leq x \leq y \text{ or } y \leq x \leq 0;$$

(iv) there exist $\delta > 0$, $\theta \in L^\infty(\Omega)$ and $\hat{\lambda} > 0$ such that

$$0 \leq \theta(z) \text{ for a.a. } z \in \Omega, \theta \not\equiv 0, \hat{\lambda} \leq \hat{\lambda}_2(q),$$

$$\theta(z)|x|^q \leq qF(z, x) \leq \hat{\lambda}|x|^q \text{ for a.a. } z \in \Omega \text{ and all } |x| \leq \delta.$$

Remark 3.1. *Evidently, hypotheses H_1 (ii), (iii) imply that for a.a. $z \in \Omega$, the function $f(z, \cdot)$ is superlinear. However, to express this superlinearity, we do not invoke the usual AR-condition. We recall that the AR-condition says that there exist $\tau > p$ and $M > 0$ such that*

$$(5) \quad 0 < \tau F(z, x) \leq f(z, x)x \text{ for a.a. } z \in \Omega \text{ and all } |x| \geq M; \text{ and}$$

$$(6) \quad 0 < \operatorname{ess\,inf}_\Omega F(\cdot, \pm M).$$

Integrating (5) and using (6), we obtain a weaker condition, namely

$$\begin{aligned} c_2|x|^\tau &\leq F(z, x) && \text{for a.a. } z \in \Omega, \text{ all } |x| \geq M \text{ and some } c_2 > 0, \\ \Rightarrow c_3|x|^\tau &\leq f(z, x)x && \text{for a.a. } z \in \Omega, \text{ all } |x| \geq M \text{ and with } c_3 = \tau c_2 > 0. \end{aligned}$$

Therefore the AR-condition implies that, eventually, $f(z, \cdot)$ has at least $(\tau-1)$ -polynomial growth.

In the present work, instead of the AR-condition, we use the quasimonotonicity hypothesis $H_1(iii)$, which is less restrictive and incorporates in our framework also $(p-1)$ -superlinear nonlinearities with slower growth near $\pm\infty$ (see the examples below). Hypothesis $H_1(iii)$ is a slight generalization of a condition which can be found in Li and Yang [11]. There are very natural ways to verify the quasimonotonicity condition. So, if there exists $M > 0$ such that for a.a. $z \in \Omega$, either the function

$$x \mapsto \frac{f(z, x)}{|x|^{q-2}x} \text{ is increasing on } x \geq M \text{ and decreasing on } x \leq -M$$

or the mapping

$$x \mapsto \eta(z, x) \text{ is increasing on } x \geq M \text{ and decreasing on } x \leq -M,$$

then hypothesis $H_1(iii)$ holds.

Hypothesis $H_1(iv)$ implies that for a.a. $z \in \Omega$, the primitive $F(z, \cdot)$ is q -linear near 0.

Examples. The following functions satisfy hypotheses H_1 . For the sake of simplicity we drop the z -dependence:

$$f_1(x) = \begin{cases} \mu|x|^{q-2}x & \text{if } |x| \leq 1 \\ \mu|x|^{r-2}x & \text{if } |x| > 1 \end{cases} \quad (\text{with } 0 < \mu \leq \hat{\lambda}_2(q) \text{ and } p < r < q^*)$$

$$f_2(x) = \begin{cases} \mu|x|^{q-2}x & \text{if } |x| \leq 1 \\ \mu|x|^{p-2}x \ln x + \mu|x|^{r-2}x & \text{if } |x| > 1 \end{cases} \quad (\text{with } 0 < \mu \leq \hat{\lambda}_2(q) \text{ and } 1 < r < p).$$

Note that only f_1 satisfies the AR-condition, whereas the function f_2 does not satisfy this growth condition.

The energy functional for problem (1) is the C^1 -functional $\varphi : W^{1,\theta}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\varphi(u) = \frac{1}{p} \gamma_p(u) + \frac{1}{q} \|Du\|_q^q - \int_{\Omega} F(z, u) dz \text{ for all } u \in W^{1,\theta}(\Omega).$$

Next, we show that $\varphi(\cdot)$ satisfies the C-condition.

Proposition 3.1. *If hypotheses H_0, H_1 hold, then the functional $\varphi(\cdot)$ satisfies the C-condition.*

Proof. We consider a sequence $\{u_n\}_{n \geq 1} \subseteq W^{1,\theta}(\Omega)$ such that

$$(7) \quad |\varphi(u_n)| \leq c_4 \text{ for some } c_4 > 0 \text{ and all } n \in \mathbb{N},$$

$$(8) \quad (1 + \|u_n\|)\varphi'(u_n) \rightarrow 0 \text{ in } W^{1,\theta}(\Omega)^* \text{ as } n \rightarrow \infty.$$

From (8) we have

$$(9) \quad \left| \langle A_p^{\alpha_0}(u_n), h \rangle + \langle A_q(u_n), h \rangle + \int_{\Omega} \xi(z)|u_n|^{p-2}u_n h dz + \int_{\partial\Omega} \beta(z)|u_n|^{p-2}u_n h d\sigma - \int_{\Omega} f(z, u_n) h dz \right| \leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|},$$

for all $h \in W^{1,\theta}(\Omega)$, with $\varepsilon_n \rightarrow 0$.

In (9) we choose $h = u_n \in W^{1,\theta}(\Omega)$ and obtain for all $n \in \mathbb{N}$

$$(10) \quad - \int_{\Omega} a_0(z)|Du_n|^p dz - \|Du_n\|_q^q - \int_{\Omega} \xi(z)|u_n|^p dz - \int_{\partial\Omega} \beta(z)|u_n|^p d\sigma + \int_{\Omega} f(z, u_n)u_n dz \leq \varepsilon_n.$$

Also, by (7) we have for all $n \in \mathbb{N}$,

$$(11) \quad \int_{\Omega} a_0(z)|Du_n|^p dz + \frac{p}{q} \|Du_n\|_q^q + \frac{p}{q} \int_{\Omega} \xi(z)|u_n|^p dz + \frac{p}{q} \int_{\partial\Omega} \beta(z)|u_n|^p d\sigma - \int_{\Omega} pF(z, u_n) dz \leq pc_4.$$

We add relations (10) and (11). Since $q < p$, we obtain

$$(12) \quad \int_{\Omega} \eta(z, u_n) dz \leq c_5 \text{ for some } c_5 > 0 \text{ and all } n \in \mathbb{N}.$$

Claim. The sequence $\{u_n\}_{n \geq 1} \subseteq W^{1,\theta}(\Omega)$ is bounded.

We argue by contradiction. Suppose that the claim is not true. We may assume that

$$(13) \quad \|u_n\| \rightarrow \infty \text{ as } n \rightarrow \infty.$$

We set $y_n = u_n/\|u_n\|$ for all $n \in \mathbb{N}$. Then $\|y_n\| = 1$ and so we may assume that

$$(14) \quad y_n \xrightarrow{w} y \text{ in } W^{1,\theta}(\Omega) \text{ and } y_n \rightarrow y \text{ in } L^r(\Omega) \text{ and in } L^p(\partial\Omega),$$

see hypotheses H_0 , Proposition 2.1 and Remark 2.1.

We first assume that $y \neq 0$. Let

$$\Omega_+ = \{z \in \Omega : y(z) > 0\} \text{ and } \Omega_- = \{z \in \Omega : y(z) < 0\}.$$

Then at least one of these measurable sets has positive Lebesgue measure on \mathbb{R}^N . We have

$$u_n(z) \rightarrow +\infty \text{ for a.a. } z \in \Omega_+ \text{ and } u_n(z) \rightarrow -\infty \text{ for a.a. } z \in \Omega_-.$$

Let $\hat{\Omega} = \Omega_+ \cup \Omega_-$ and let $|\cdot|_N$ denote the Lebesgue measure on \mathbb{R}^N . We see that $|\hat{\Omega}|_N > 0$ and on account of hypothesis $H_1(ii)$, we have

$$(15) \quad \begin{aligned} \frac{F(z, u_n(z))}{\|u_n\|^p} &= \frac{F(z, u_n(z))}{|u_n(z)|^p} |y_n(z)|^p \rightarrow +\infty \text{ for a.a. } z \in \hat{\Omega}, \\ \Rightarrow \int_{\hat{\Omega}} \frac{F(z, u_n(z))}{\|u_n\|^p} dz &\rightarrow +\infty \text{ by Fatou's lemma.} \end{aligned}$$

Hypotheses $H_1(i)$, (ii) imply that

$$(16) \quad F(z, x) \geq -c_6 \text{ for a.a. } z \in \Omega, \text{ all } x \in \mathbb{R} \text{ and some } c_6 > 0.$$

Thus we obtain

$$(17) \quad \begin{aligned} \int_{\Omega} \frac{F(z, u_n)}{\|u_n\|^p} dz &= \int_{\hat{\Omega}} \frac{F(z, u_n)}{\|u_n\|^p} dz + \int_{\Omega \setminus \hat{\Omega}} \frac{F(z, u_n)}{\|u_n\|^p} dz \\ &\geq \int_{\hat{\Omega}} \frac{F(z, u_n)}{\|u_n\|^p} dz - \frac{c_6 |\Omega|_N}{\|u_n\|^p} \text{ (see (16)),} \\ &\Rightarrow \lim_{n \rightarrow \infty} \int_{\Omega} \frac{F(z, u_n)}{\|u_n\|^p} dz = +\infty \text{ (see (15) and (13)).} \end{aligned}$$

By (7), we have

$$(18) \quad \int_{\Omega} \frac{pF(z, u_n)}{\|u_n\|^p} dz \leq \gamma_p(y_n) + \frac{p}{q} \frac{1}{\|u_n\|^{p-q}} \|Dy_n\|_q^q + \frac{c_4}{\|u_n\|^p} \leq c_7,$$

for some $c_7 > 0$ and all $n \in \mathbb{N}$ (see (13) and recall that $\|y_n\| = 1$).

We compare relations (15) and (18) and arrive at a contradiction.

Next, we assume that $y = 0$. Let $\mu > 0$ and set $v_n = (p\mu)^{1/p} y_n$ for all $n \in \mathbb{N}$. Evidently, we have

$$(19) \quad \begin{aligned} v_n &\rightarrow 0 \text{ in } L^r(\Omega) \text{ (see (14)),} \\ \Rightarrow \int_{\Omega} F(z, v_n) dz &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Consider the functional $\psi : W^{1,\theta}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\psi(u) = \frac{1}{p} \gamma_p(u) - \int_{\Omega} F(z, u) dz \text{ for all } u \in W^{1,\theta}(\Omega).$$

Clearly, $\psi \in C^1(W^{1,\theta}(\Omega), \mathbb{R})$ and

$$(20) \quad \psi \leq \varphi.$$

We can find $t_n \in [0, 1]$ such that

$$(21) \quad \psi(t_n u_n) = \min\{\psi(tu_n) : 0 \leq t \leq 1\} \text{ for all } n \in \mathbb{N}.$$

Because of (13), we can find $n_0 \in \mathbb{N}$ such that

$$(22) \quad 0 < \frac{(p\mu)^{1/p}}{\|u_n\|} \leq 1 \text{ for all } n \geq n_0.$$

Therefore

$$\begin{aligned}
\psi(t_n u_n) &\geq \psi(v_n) \text{ (see (21), (22))} \\
&\geq \mu \gamma_p(y_n) - \int_{\Omega} F(z, v_n) dz \\
&\geq \mu c_1 - \int_{\Omega} F(z, v_n) dz \text{ (see (2) and recall that } \|y_n\| = 1) \\
&\geq \frac{\mu}{2} c_1 \text{ for all } n \geq n_1 \geq n_0 \text{ (see (19)).}
\end{aligned}$$

Since $\mu > 0$ is arbitrary, it follows that

$$(23) \quad \psi(t_n u_n) \rightarrow +\infty \text{ as } n \rightarrow \infty.$$

Note that

$$(24) \quad \psi(0) = 0 \text{ and } \psi(u_n) \leq c_4 \text{ for all } n \in \mathbb{N} \text{ (see (7), (20)).}$$

By (23) and (24) we can infer that

$$(25) \quad t_n \in (0, 1) \text{ for all } n \geq n_2.$$

From (21) and (25), we can see that for all $n \geq n_2$ we have

$$\begin{aligned}
(26) \quad 0 &= t_n \frac{d}{dt} \psi(tu_n)|_{t=t_n} \\
&= \langle \psi'(t_n u_n), t_n u_n \rangle \text{ (by the chain rule)} \\
&= \gamma_p(t_n u_n) - \int_{\Omega} f(z, t_n u_n)(t_n u_n) dz.
\end{aligned}$$

It follows that

$$0 \leq t_n u_n^+ \leq u_n^+ \text{ and } -u_n^- \leq -t_n u_n^- \leq 0 \text{ for all } n \in \mathbb{N}$$

(recall that $u_n^+ = \max\{u_n, 0\}$ and $u_n^- = \max\{-u_n, 0\}$).

By hypothesis $H_1(iii)$, we have

$$\eta(z, t_n u_n^+) \leq \eta(z, u_n^+) + e(z) \text{ for a.a. } z \in \Omega \text{ and all } n \in \mathbb{N},$$

$$\eta(z, -t_n u_n^-) \leq \eta(z, -u_n^-) + e(z) \text{ for a.a. } z \in \Omega \text{ and all } n \in \mathbb{N}.$$

From these two inequalities and since $u_n = u_n^+ - u_n^-$, we obtain

$$(27) \quad \begin{aligned} &\eta(z, t_n u_n) \leq \eta(z, u_n) + e(z) \text{ for a.a. } z \in \Omega \text{ and all } n \in \mathbb{N}, \\ \Rightarrow &f(z, t_n u_n)(t_n u_n) \leq \eta(z, u_n) + e(z) + pF(z, t_n u_n) \text{ for a.a. } z \in \Omega \text{ and all } n \in \mathbb{N}. \end{aligned}$$

We return to (26) and apply (27). Then

$$(28) \quad \begin{aligned} &\gamma_p(t_n u_n) - p \int_{\Omega} F(z, t_n u_n) dz \leq \int_{\Omega} \eta(z, u_n) dz + \|e\|_1 \text{ for all } n \in \mathbb{N}, \\ \Rightarrow &p\psi(t_n u_n) \leq c_8 \text{ for some } c_8 > 0 \text{ and all } n \in \mathbb{N} \text{ (see (12)).} \end{aligned}$$

We compare (23) and (28) and arrive at a contradiction.

This proves the claim.

On account of this claim, we may assume that

$$(29) \quad u_n \xrightarrow{w} u \text{ in } W^{1,\theta}(\Omega) \text{ and } u_n \rightarrow u \text{ in } L^r(\Omega) \text{ and in } L^p(\partial\Omega)$$

(see hypotheses H_0).

From (29) we have

$$(30) \quad Du_n \rightarrow Du \text{ in } L^p_{a_0}(\Omega, \mathbb{R}^N) \quad \text{and} \quad Du_n(z) \rightarrow Du(z) \text{ a.a. } z \in \Omega.$$

In (9) we choose $h = u_n - u \in W^{1,\theta}(\Omega)$, pass to the limit as $n \rightarrow \infty$ and use (30) and the monotonicity of $A_p(\cdot)^{a_0}$. We obtain

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \langle A_p^{a_0}(u_n), u_n - u \rangle \leq 0, \\ \Rightarrow &\limsup_{n \rightarrow \infty} \|Du_n\|_{L^p_{a_0}(\Omega, \mathbb{R}^N)} \leq \|Du\|_{L^p_{a_0}(\Omega, \mathbb{R}^N)}. \end{aligned}$$

On the other hand, from (30) we have

$$\liminf_{n \rightarrow \infty} \|Du_n\|_{L_{a_0}^p(\Omega, \mathbb{R}^N)} \geq \|Du\|_{L_{a_0}^p(\Omega, \mathbb{R}^N)}.$$

Therefore we conclude that

$$(31) \quad \|Du_n\|_{L_{a_0}^p(\Omega, \mathbb{R}^N)} \rightarrow \|Du\|_{L_{a_0}^p(\Omega, \mathbb{R}^N)}.$$

The space $L_{a_0}^p(\Omega, \mathbb{R}^N)$ is uniformly convex, hence it has the Kadec-Klee property (see Papa-georgiou, Rădulescu and Repovš [18, Remark 2.7.30, p. 127]). So, it follows from (30) and (31) that

$$\begin{aligned} & Du_n \rightarrow Du \text{ in } L_{a_0}^p(\Omega, \mathbb{R}^N), \\ \Rightarrow & Du_n \rightarrow Du \text{ in } L^q(\Omega, \mathbb{R}^N) \text{ since } L_{a_0}^p(\Omega, \mathbb{R}^N) \hookrightarrow L^q(\Omega, \mathbb{R}^N) \text{ continuously,} \\ \Rightarrow & \rho_\theta(|Du_n - Du|) \rightarrow 0 \text{ (see Proposition 2.2),} \\ \Rightarrow & \|u_n - u\| \rightarrow 0 \text{ (see (29) and Proposition 2.2),} \\ \Rightarrow & \varphi \text{ satisfies the C-condition.} \end{aligned}$$

The proof is now complete. \square

Proposition 3.2. *If hypotheses H_0, H_1 hold, then the functional $\varphi(\cdot)$ has a local $(1, 1)$ -linking at 0.*

Proof. Since the critical points of φ are solutions of problem (1), we may assume that K_φ is finite or otherwise we already have infinitely many nontrivial solutions of (1) and so we are done.

Choose $\rho \in (0, 1)$ so small that $K_\varphi \cap \bar{B}_\rho = \{0\}$ (here, $B_\rho = \{u \in W^{1,\theta}(\Omega) : \|u\| < \rho\}$). Let $V = \mathbb{R}$ and let $\delta > 0$ as postulated by hypothesis $H_1(iv)$. Recall that on a finite dimensional normed space all norms are equivalent. So, by taking $\rho \in (0, 1)$ even smaller as necessary, we have

$$(32) \quad \|u\| \leq \rho \Rightarrow |u| \leq \delta \text{ for all } u \in V = \mathbb{R}.$$

Then for $u \in V \cap \bar{B}_\rho$, we have

$$\begin{aligned} \varphi(u) & \leq \frac{1}{p} \gamma_p(u) - \frac{|u|^q}{q} \int_\Omega \theta(z) dz \text{ (see (32) and Hypothesis } H_1(iv)) \\ & = \frac{|u|^p}{p} \left(\int_\Omega \xi(z) dz + \int_{\partial\Omega} \beta(z) d\sigma \right) - \frac{|u|^q}{q} \int_\Omega \theta(z) dz \\ & \leq c_9 \|u\|^p - c_{10} \|u\|^q \text{ for some } c_9, c_{10} > 0 \text{ (see hypotheses } H_0 \text{ and } H_1(iv)). \end{aligned}$$

Since $q < p$, choosing $\rho \in (0, 1)$ small, we conclude that

$$(33) \quad \varphi|_{V \cap \bar{B}_\rho} \leq 0.$$

Let

$$D = \{u \in W^{1,\theta}(\Omega) : \|Du\|_q^q \geq \hat{\lambda}_2(q) \|u\|_q^q\}.$$

For all $u \in D$ we have

$$\begin{aligned} \varphi(u) & = \frac{1}{p} \gamma_p(u) + \frac{1}{q} \|Du\|_q^q - \int_{\{|u| \leq \delta\}} F(z, u) dz - \int_{\{|u| > \delta\}} F(z, u) dz \\ & \geq \frac{1}{p} \gamma_p(u) + \frac{1}{q} \left(\|Du\|_q^q - \int_\Omega \hat{\lambda} |u|^q dz \right) - \int_\Omega F(z, u) dz \\ & \quad \text{(see hypotheses } H_1(iv)) \\ & \geq \frac{1}{p} \gamma_p(u) + \frac{1}{q} \int_\Omega (\hat{\lambda}_2(q) - \hat{\lambda}) |u|^q dz - c_{11} \|u\|^r \\ & \quad \text{for some } c_{11} > 0 \text{ (since } u \in D \text{ and see hypothesis } H_1(iv)) \\ & \geq \frac{c_{11}}{p} \|u\|^p - c_{11} \|u\|^r \text{ (see (22)).} \end{aligned}$$

Since $p < r$, for small $\rho \in (0, 1)$ we have

$$(34) \quad \varphi|_{D \cap \bar{B}_\rho \setminus \{0\}} > 0.$$

Let $U = \bar{B}_\rho$, $E_0 = V \cap \partial B_\rho$, $E = V \cap \bar{B}_\rho$ and D as above. We have $0 \notin E_0$, $E_0 \subseteq E \subseteq U = \bar{B}_\rho$ and $E_0 \cap D = \emptyset$ (see Definition 2.3).

Let Y be the topological complement of V . We have that

$$W^{1,\theta}(\Omega) = V \oplus Y \text{ (see [18, pp. 73, 74]).}$$

So, every $u \in W^{1,\theta}(\Omega)$ can be written in a unique way as

$$u = v + y \text{ with } v \in V, y \in Y.$$

We consider the deformation $h : [0, 1] \times (W^{1,\theta}(\Omega) \setminus D) \rightarrow W^{1,\theta}(\Omega) \setminus D$ defined by

$$h(t, u) = (1-t)u + t\rho \frac{v}{\|v\|} \text{ for all } t \in [0, 1], u \in W^{1,\theta}(\Omega) \setminus D.$$

We have

$$h(0, u) = u \text{ and } h(1, u) = \rho \frac{v}{\|v\|} \in V \cap \partial B_\rho = E_0.$$

It follows that E_0 is a deformation retract of $W^{1,\theta}(\Omega) \setminus D$ (see Papageorgiou, Rădulescu and Repovš [17, Definition 5.3.10, p. 385]). Hence

$$i_* : H_0(E_0) \rightarrow H_0(W^{1,\theta}(\Omega) \setminus \{0\})$$

is an isomorphism (see Eilenberg and Steenrod [8, Theorem 11.5, p.30] and Papageorgiou, Rădulescu and Repovš [18, Remark 6.1.6, p. 460]).

The set $E = V \cap B_\rho$ is contractible (it is an interval). Hence $H_0(E, E_0) = 0$ (see Eilenberg and Steenrod [8, Theorem 11.5, p. 30]). Therefore, if $j_* : H_0(E_0) \rightarrow H_0(E)$, then $\dim \text{im } j_* = 1$ (see Papageorgiou, Rădulescu and Repovš [8, Remark 6.1.26, p. 468]). So, finally we have

$$\begin{aligned} & \dim \text{im } i_* - \dim \text{im } j_* = 2 - 1 = 1, \\ \Rightarrow & \varphi(\cdot) \text{ has a local } (1, 1)\text{-linking at } 0, \text{ see Definition 2.3.} \end{aligned}$$

The proof is now complete. \square

From Proposition 3.2 and Theorem 6.6.17 of Papageorgiou, Rădulescu and Repovš [18, p. 538], we have

$$(35) \quad \dim C_1(\varphi, 0) \geq 1.$$

Moreover, Proposition 3.9 of Papageorgiou, Rădulescu and Repovš [17] leads to the following result.

Proposition 3.3. *If hypotheses H_0, H_1 hold, then $C_k(\varphi, \infty) = 0$ for all $k \in \mathbb{N}_0$.*

We are now ready for the existence theorem concerning the superlinear case.

Theorem 3.4. *If hypotheses H_0, H_1 hold, then problem (1) has a nontrivial solution $u_0 \in W^{1,\theta}(\Omega) \cap L^\infty(\Omega)$.*

Proof. On account of (35) and Proposition 3.3, we can apply Proposition 6.2.42 of Papageorgiou, Rădulescu and Repovš [18, p. 499]. So, we can find $u_0 \in W^{1,\theta}(\Omega)$ such that

$$\begin{aligned} & u_0 \in K_\varphi \setminus \{0\}, \\ \Rightarrow & u_0 \in W^{1,\theta}(\Omega) \cap L^\infty(\Omega) \text{ is a solution of problem (1), see [18, Section 3.2].} \end{aligned}$$

The proof is now complete. \square

4. RESONANT CASE

In this section we are concerned with the resonant case (p -linear case). Our hypotheses allow resonance at $\pm\infty$ with respect to the principal eigenvalue $\hat{\lambda}_1(p) > 0$.

The new conditions on the reaction $f(z, x)$ are the following.

- H_2 : $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function such that $f(z, 0) = 0$ for a.a. $z \in \Omega$ and
- (i) $|f(z, x)| \leq \hat{a}(z)(1 + |x|^{r-1})$ for a.a. $z \in \Omega$, all $x \in \mathbb{R}$, with $\hat{a} \in L^\infty(\Omega)$, $p < r < q^*$;
 - (ii) if $F(z, x) = \int_0^x f(z, s)ds$, then $\lim_{x \rightarrow \pm\infty} pF(z, x)/|x|^p \leq \hat{\lambda}_1(p)$ uniformly for a.a. $z \in \Omega$;
 - (iii) we have

$$f(z, x)x - pF(z, x) \rightarrow +\infty \text{ uniformly for a.a. } z \in \Omega, \text{ as } x \rightarrow \pm\infty;$$

- (iv) there exist $\delta > 0$, $\theta \in L^\infty(\Omega)$ and $\hat{\lambda} > 0$ such that

$$\begin{aligned} & 0 \leq \theta(z) \text{ for a.a. } z \in \Omega, \theta \not\equiv 0, \hat{\lambda} \leq \hat{\lambda}_2(q), \\ & \theta(z)|x|^q \leq qF(z, x) \leq \hat{\lambda}|x|^q \text{ for a.a. } z \in \Omega \text{ and all } |x| \leq \delta. \end{aligned}$$

Remark 4.1. Hypothesis $H_2(ii)$ implies that at $\pm\infty$, we can have resonance with respect to the principal eigenvalue of the operator $u \mapsto -\operatorname{div}(a_0(z)|Du|^{p-2}Du) - \Delta_q u$ with Robin boundary condition.

Proposition 4.1. If hypotheses H_0, H_2 hold, then the energy functional $\varphi(\cdot)$ is coercive.

Proof. We have

$$\begin{aligned} \frac{d}{dx} \left(\frac{F(z, x)}{|x|^p} \right) &= \frac{f(z, x)|x|^p - p|x|^{p-2}xF(z, x)}{|x|^{2p}} \\ &= \frac{|x|^{p-2}x[f(z, x)x - pF(z, x)]}{|x|^{2p}}. \end{aligned}$$

On account of hypothesis $H_2(iii)$, given any $\gamma > 0$, we can find $M_1 = M_1(\gamma) > 0$ such that

$$f(z, x)x - pF(z, x) \geq \gamma \text{ for a.a. } z \in \Omega \text{ and all } |x| \geq M_1.$$

Hence we obtain

$$\frac{d}{dx} \left(\frac{F(z, x)}{|x|^p} \right) \begin{cases} \geq \frac{\gamma}{x^{p+1}} & \text{if } x \geq M_1 \\ \leq -\frac{\gamma}{|x|^{p+1}} & \text{if } x \leq -M_1. \end{cases}$$

Integrating, we obtain

$$(36) \quad \frac{F(z, x)}{|x|^p} - \frac{F(z, u)}{|u|^p} \geq -\frac{\gamma}{p} \left(\frac{1}{|x|^p} - \frac{1}{|u|^p} \right) \text{ for a.a. } z \in \Omega \text{ and all } |x| \geq |u| \geq M_1.$$

On account of hypothesis $H_2(ii)$, given $\varepsilon > 0$, we can find $M_2 = M_2(\varepsilon) > 0$ such that

$$F(z, x) \leq \frac{1}{p} (\hat{\lambda}_1(p) + \varepsilon) |x|^p \text{ for a.a. } z \in \Omega \text{ and all } |x| \geq M_2.$$

Using this inequality in (36) and letting $|x| \rightarrow \infty$ we obtain

$$(37) \quad \begin{aligned} \frac{1}{p} (\hat{\lambda}_1(p) + \varepsilon) - \frac{F(z, u)}{|u|^p} &\geq \frac{\gamma}{p} \frac{1}{|u|^p} \text{ for a.a. } z \in \Omega \text{ and all } |u| \geq M = \max\{M_1, M_2\}, \\ \Rightarrow (\hat{\lambda}_1(p) + \varepsilon) |u|^p - pF(z, u) &\geq \gamma \text{ for a.a. } z \in \Omega \text{ and all } |u| \geq M. \end{aligned}$$

Arguing by contradiction, suppose that $\varphi(\cdot)$ is not coercive. Then we can find $\{u_n\}_{n \geq 1} \subseteq W^{1, \theta}(\Omega)$ such that

$$(38) \quad \|u_n\| \rightarrow \infty \text{ and } \varphi(u_n) \leq M_0 \text{ for some } M_0 > 0 \text{ and all } n \in \mathbb{N}.$$

Let $y_n = u_n / \|u_n\|$ for all $n \in \mathbb{N}$. Then $\|y_n\| = 1$, hence we may assume that

$$(39) \quad y_n \xrightarrow{w} y \text{ in } W^{1, \theta}(\Omega) \text{ and } y_n \rightarrow y \text{ in } L^p(\Omega) \text{ and in } L^p(\partial\Omega).$$

From (38) we have

$$\begin{aligned} &\frac{1}{p} \gamma_p(y_n) + \frac{1}{q} \frac{1}{\|u_n\|^{p-q}} \int_{\Omega} |Dy_n|^q dz - \int_{\Omega} \frac{F(z, u_n)}{\|u_n\|^p} dz \leq \frac{M_0}{\|u_n\|^p}, \\ \Rightarrow \gamma_p(y_n) + \frac{p}{q} \frac{1}{\|u_n\|^{p-q}} \int_{\Omega} |Dy_n|^q dz &\leq \tau_n + (\hat{\lambda}_1(p) + \varepsilon) \|y_n\|_p^p \text{ with } \tau_n \rightarrow 0, \text{ see (37),} \\ \Rightarrow \gamma_p(y) &\leq (\hat{\lambda}_1(p) + \varepsilon) \|y\|_p^p \text{ (see (39)),} \\ \Rightarrow \gamma_p(y) &\leq \hat{\lambda}_1(p) \|y\|_p^p \text{ (since } \varepsilon > 0 \text{ is arbitrary),} \\ \Rightarrow y &= \mu \hat{u}_1(p) \text{ for some } \mu \in \mathbb{R} \text{ (see (4)).} \end{aligned}$$

If $\mu = 0$, then $y = 0$ and so $\gamma_p(y_n) \rightarrow 0$. Hence, as in the proof of Proposition 3.1, we have $y_n \rightarrow 0$ in $W^{1, \theta}(\Omega)$, contradicting the fact that $\|y_n\| = 1$ for all $n \in \mathbb{N}$.

So, $\mu \neq 0$ and since $\hat{u}_1(p)(z) > 0$ for a.a. $z \in \Omega$, we have $|u_n(z)| \rightarrow +\infty$ for a.a. $z \in \Omega$. By (38) and (4) we have

$$(40) \quad \int_{\Omega} \left[\frac{1}{p} \hat{\lambda}_1(p) |u_n|^p - F(z, u_n) \right] dz \leq M_0 \text{ for all } n \in \mathbb{N}.$$

However, from (37) and since $\gamma > 0$ is arbitrary, we can infer that

$$(41) \quad \begin{aligned} & \frac{1}{p} \hat{\lambda}_1(p) |u_n|^p - F(z, u_n) \rightarrow +\infty \text{ for a.a. } z \in \Omega, \text{ as } n \rightarrow \infty, \\ \Rightarrow & \int_{\Omega} \left[\frac{1}{p} \hat{\lambda}_1(p) |u_n|^p - F(z, u_n) \right] dz \rightarrow +\infty \text{ by Fatou's lemma.} \end{aligned}$$

Comparing (40) and (41) we arrive at a contradiction. Therefore we can conclude that $\varphi(\cdot)$ is coercive. \square

Using Proposition 4.1 and Proposition 5.1.15 of Papageorgiou, Rădulescu and Repovš [18, p. 369], we obtain the following result.

Corollary 4.2. *If hypotheses H_0, H_2 hold, then the energy functional $\varphi(\cdot)$ is bounded below and satisfies the C-condition.*

Now we are ready for the multiplicity theorem in the resonant case.

Theorem 4.3. *If hypotheses H_0, H_2 hold, then problem (1) has at least two nontrivial solutions $u_0, \hat{u} \in W^{1,\theta}(\Omega) \cap L^\infty(\Omega)$.*

Proof. By Proposition 3.2 we know that $\varphi(\cdot)$ has a local (1,1)-linking at the origin. Note that for that result mattered only the behavior of $f(z, \cdot)$ near zero and this is common in hypotheses H_1 and H_2 . Also, we know that $\varphi(\cdot)$ is sequentially weakly lower semicontinuous. This fact in conjunction with Proposition 4.1, permit the use of the Weierstrass-Tonelli theorem. So, we can find $u_0 \in W^{1,\theta}(\Omega)$ such that

$$(42) \quad \varphi(u_0) = \min\{\varphi(u) : u \in W^{1,\theta}(\Omega)\}.$$

On account of hypothesis $H_2(iv)$ and since $q < p$, we have

$$\begin{aligned} & \varphi(u_0) < 0 = \varphi(0), \\ \Rightarrow & u_0 \neq 0 \text{ and } u_0 \in K_\varphi, \\ \Rightarrow & u_0 \in K_\varphi \cap L^\infty(\Omega) \text{ is a nontrivial solution of (1).} \end{aligned}$$

Moreover, by Corollary 6.7.10 of Papageorgiou, Rădulescu and Repovš [18, p. 552], we can find $\hat{u} \in K_\varphi, \hat{u} \notin \{0, u_0\}$. Then $\hat{u} \in W^{1,\theta}(\Omega) \cap L^\infty(\Omega)$ is the second nontrivial solution of problem (1). \square

Acknowledgments. This research was supported by the Slovenian Research Agency grants P1-0292, J1-8131, N1-0114, N1-0064, and N1-0083.

REFERENCES

- [1] A. Bahrouni, V.D. Rădulescu, D.D. Repovš, A weighted anisotropic variant of the Caffarelli-Kohn-Nirenberg inequality and applications, *Nonlinearity* **31** (2018), no. 4, 1516-1534.
- [2] A. Bahrouni, V.D. Rădulescu, D.D. Repovš, Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves, *Nonlinearity* **32** (2019), no. 7, 2481-2495.
- [3] P. Baroni, M. Colombo, G. Mingione, Harnack inequalities for double phase functionals, *Nonlinear Anal.* **121** (2015), 206-222.
- [4] M. Cencelj, V.D. Rădulescu, D.D. Repovš, Double phase problems with variable growth, *Nonlinear Anal.* **177** (2018), part A, 270-287.
- [5] F. Colasuonno, M. Squassina, Eigenvalues for double phase variational integrals, *Ann. Mat. Pura Appl.* (4) **195** (2016), no. 6, 1917-1959.
- [6] M. Colombo, G. Mingione, Regularity for double phase variational problems, *Arch. Ration. Mech. Anal.* **215** (2015), no. 2, 443-496.
- [7] M. Colombo, G. Mingione, Bounded minimisers of double phase variational integrals, *Arch. Ration. Mech. Anal.* **218** (2015), no. 1, 219-273.
- [8] S. Eilenberg, N. Steenrod, *Foundations of Algebraic Topology*, Princeton University Press, Princeton, New Jersey, 1952.
- [9] L. Gasinski, N.S. Papageorgiou, *Nonlinear Analysis*, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [10] L. Gasinski, N.S. Papageorgiou, Positive solutions for the Robin p -Laplacian problem with competing nonlinearities, *Adv. Calc. Var.* **12** (2019), no. 1, 31-56.
- [11] G. Li, C. Yang, The existence of a nontrivial solution to a nonlinear elliptic boundary value problem of p -Laplacian type without the Ambrosetti-Rabinowitz condition, *Nonlinear Anal.* **72** (2010), no. 12, 4602-4613.

- [12] J. Liu, The Morse index of a saddle point, *Systems Sci. Math. Sci.* **2** (1989), no. 1, 32-39.
- [13] W. Liu, G. Dai, Existence and multiplicity results for double phase problem, *J. Differential Equations* **265** (2018), no. 9, 4311-4334.
- [14] P. Marcellini, Regularity and existence of solutions of elliptic equations with p, q -growth conditions, *J. Differential Equations* **90** (1991), no. 1, 1-30.
- [15] D. Mugnai, N.S. Papageorgiou, Resonant nonlinear Neumann problems with indefinite weight, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **11** (2012), no. 4, 729-788.
- [16] J. Musielak, *Orlicz Spaces and Modular Spaces*, Lecture Notes in Mathematics, vol. 1034, Springer-Verlag, Berlin, 1983.
- [17] N.S. Papageorgiou, V.D. Rădulescu, Nonlinear nonhomogeneous Robin problems with superlinear reaction term, *Adv. Nonlinear Stud.* **16** (2016), no. 4, 737-764.
- [18] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, *Nonlinear Analysis—Theory and Methods*, Springer Monographs in Mathematics, Springer Nature, Cham, 2019.
- [19] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, Ground state and nodal solutions for a class of double phase problems, *Z. Angew. Math. Phys.* **71** (2020), no. 1, art. 15, 15 pp.
- [20] N.S. Papageorgiou, C. Vetro, F. Vetro, Multiple solutions for parametric double phase Dirichlet problems, *Commun. Contemp. Math.*, in press (<https://doi.org/10.1142/S0219199720500066>).
- [21] K. Perera, Homological local linking, *Abstr. Appl. Anal.* **3** (1998), no. 1-2, 181-189.
- [22] V.V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory (Russian); English transl. in *Izv. Akad. Nauk SSSR Ser. Mat.* **50** (1986), no. 4, 675-710, 877.

(N.S. Papageorgiou) INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, 1000 LJUBLJANA, SLOVENIA & DEPARTMENT OF MATHEMATICS, NATIONAL TECHNICAL UNIVERSITY, ZOGRAFOU CAMPUS, ATHENS 15780, GREECE
Email address: `npapg@math.ntua.gr`

(V.D. Rădulescu) INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, 1000 LJUBLJANA, SLOVENIA & FACULTY OF APPLIED MATHEMATICS, AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, 30-059 KRAKÓW, POLAND
Email address: `vicentiu.radulescu@imfm.si`

(D.D. Repovš) FACULTY OF EDUCATION AND FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA & INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, 1000 LJUBLJANA, SLOVENIA
Email address: `dusan.repovs@guest.arnes.si`